

# MODULI SPACES OF CURVES

SCOTT NOLLET

ABSTRACT. My goal is to introduce vocabulary and present examples that will help graduate students to better follow lectures at TAGS 2018. Assuming some background in modern algebraic geometry (varieties, schemes and cohomology), I will discuss linear systems on curves, Hilbert schemes, and the Mumford-Deligne moduli space of isomorphism classes of curves of fixed genus.

## 1. LINEAR SYSTEMS ON CURVES

In this section we establish the curves to be studied, the notion of linear system, and the Riemann-Roch theorem and some of its applications. We work throughout over a fixed algebraically closed field  $k$  of arbitrary characteristic.

1.1. **Curves.** We first clarify our main object of study:

**Definition 1.1.** An embedded curve  $C \subset \mathbb{P}_k^n$  is a nonsingular one dimensional closed subvariety [4, I,§5]. Letting  $P_C(t) = dt + a$  be the Hilbert polynomial, the geometric genus of  $C$  is  $g = 1 - a$  and the degree is  $d = \deg C$ .

An abstract curve  $C$  is a one dimensional nonsingular complete variety of finite type over  $k$  [4, IV,§1]. Letting  $\omega_C$  be the sheaf of differentials<sup>1</sup> [4, II,§8], the geometric genus of  $C$  is  $g = \dim H^0(\omega_C) = \dim H^1(\mathcal{O}_C)$ . If  $k = \mathbb{C}$ , then  $g$  is the number of handles on the corresponding real surface.

**Remark 1.2.** An embedded curve  $C \subset \mathbb{P}^n$  yields  $C = \text{Proj } k[x_0, \dots, x_n]/I_C$  as an abstract curve, but the embedded curve has extra data such as the ideal  $I_C$  and the degree  $\deg C$ .

**Remark 1.3.** One can also define an abstract curve as the set of all discrete valuation rings  $k \subset R \subset K$ , where  $k \subset K$  has transcendence degree one [4, I,§6].

**Example 1.4.** A general hypersurface  $C \subset \mathbb{P}^2$  of degree  $d$  is a curve with  $g = \binom{d-1}{2}$ .

**Example 1.5.** [4, III,Exercise 5.6] For  $a, b > 0$ , general effective divisors on the nonsingular quadric  $Q \subset \mathbb{P}^3$  of type  $\mathcal{O}_Q(a, b)$  are curves  $C$  with  $g = (a-1)(b-1)$  and  $\deg C = a+b$ . Taking  $a = 2, b \geq 1$  gives curves of every genus  $g \geq 0$ .

1.2. **Linear systems.** Any map  $f : C \rightarrow \mathbb{P}^n$  gives rise to a line bundle  $L = f^*\mathcal{O}_{\mathbb{P}^n}(1)$  and sections  $s_i = f^*x_i$  which generate  $L$  in the sense that  $\mathcal{O}_C^{n+1} \xrightarrow{s_i} L$  is surjective.

$$\{f : C \rightarrow \mathbb{P}^n\} \longleftrightarrow \left\{ \begin{array}{l} L \in \text{Pic } C \\ s_0, \dots, s_n \in H^0(C, L) \\ s_i \text{ generate } L \end{array} \right\} \longrightarrow \left\{ \begin{array}{l} L \in \text{Pic } C \\ V = \langle s_0, \dots, s_n \rangle \in H^0(C, L) \end{array} \right\}$$

---

<sup>1</sup>Namely  $\omega_C = \Delta^*(\mathcal{I}/\mathcal{I}^2)$ , where  $\mathcal{I}$  is the ideal of  $\Delta$  in  $C \times C$ . We use  $K_C$  for the associated divisor.

Geometrically a hyperplane  $H$  with equation  $\sum a_i x_i$  pulls back to the zero set of the section  $s = \sum a_i s_i$ , denoted  $(s)_0$ : dualize the injection  $s : \mathcal{O}_C \rightarrow L$  to get  $L^\vee \hookrightarrow \mathcal{O}_C$  as the ideal sheaf of the effective divisor  $D = (s)_0 = \sum n_i p_i$  of degree  $\deg D = \sum n_i$ . These are determined by  $s$  up to  $\lambda \in k^*$ , so the set of all such is given by the projective space

$$\mathbb{P}V = V - \{0\} / \sim .$$

This is a linear system of dimension  $r = \dim V - 1$  and degree  $d = \deg L$ , denoted  $g_d^r$ .

**Example 1.6.** (a) Take  $C = \mathbb{P}^1, L = \mathcal{O}_{\mathbb{P}^1}(3)$  and  $V = H^0(\mathbb{P}^1, L)$ . For homogeneous coordinates  $s, t$  on  $\mathbb{P}^1$  and  $V = \langle s^3, s^2t, st^2, t^3 \rangle$ , the associated map  $\mathbb{P}^1 \rightarrow \mathbb{P}^3$  is given by

$$(s, t) \mapsto (x, y, z, w) = (s^3, s^2t, st^2, t^3).$$

The image  $C$  has ideal  $I_C = (xz - y^2, xw - yz, yw - z^2)$ . Modulo  $\text{Aut } \mathbb{P}^3$ , every curve of degree  $d = 3$  and genus  $g = 0$  arises from this construction [4, II, Example 7.8.5] and is called a twisted cubic curve.

Two more linear systems are present. Note that  $C \subset Q$  is a divisor of type  $(2, 1)$ , so there are two projections to  $\mathbb{P}^1$  via the isomorphism  $Q \cong \mathbb{P}^1 \times \mathbb{P}^1$ :

- (b)  $\pi_1$  is an isomorphism giving a  $g_1^1$ . A curve  $C$  carries a  $g_1^1 \iff C \cong \mathbb{P}^1$
- (c)  $\pi_2$  is a double cover giving a  $g_2^1$ .

**Exercise 1.7.** In Example 1.6 (c), compute  $V \subset H^0(\mathcal{O}_{\mathbb{P}^1}(2))$ .

**Remark 1.8.** A curve  $C$  carrying a  $g_2^1$  is a double cover of  $\mathbb{P}^1$  and is called hyperelliptic.

**1.3. Riemann-Roch.** The Riemann Roch theorem estimates the dimension  $l(D)$  of the complete linear system  $|D|$  associated to a divisor  $D = \sum n_i p_i$ . If  $H^0(C, \mathcal{O}_C(D)) = 0$ , then there are no effective divisors linearly equivalent to  $D$ , so we take  $l(D) = -1$ .

**Theorem 1.9.** *If  $D$  is a divisor on  $C$ , then  $l(D) - l(K_C - D) = \deg D + 1 - g$ .*

**Corollary 1.10.**  $\deg K_C = 2g - 2$ . [Take  $D = K_C$  so that  $l(K_C) = g - 1$  and  $l(0) = 0$ .]

**Corollary 1.11.** *If  $\deg D > 2g - 2$ , then  $l(D) = \deg D - g$ .* [Because  $l(K_C - D) = -1$ .]

**Remark 1.12.** In terms of the map, we observe the following [4, IV, Proposition 3.1].

- (a)  $P \in C$  is a base point for  $|D|$  iff the map  $|D - P| \hookrightarrow |D|$  given by  $A \mapsto A + P$  is not onto, hence  $|D|$  is base point free iff  $l(D - P) = l(D) - 1$  for each  $P$ .
- (b) Similarly  $|D|$  separates points and tangent vectors iff  $l(D - P - Q) = l(D) - 2$  for each  $P, Q \in C$ , including the case  $P = Q$ .

**Corollary 1.13.** *If  $d = \deg D \geq 2g + 1$ , then  $|D|$  gives a closed immersion  $C \hookrightarrow \mathbb{P}^{d-g}$ .*

*Proof.* For  $P, Q \in C, D, D - P$  and  $D - P - Q$  are nonspecial, so apply Corollary 1.11.  $\square$

**Example 1.14.** Three points on an elliptic curve  $C$  give a closed immersion  $C \hookrightarrow \mathbb{P}^2$ .

2. HILBERT SCHEMES

A flat family in  $\mathbb{P}^n$  consists of a closed immersion  $X \hookrightarrow \mathbb{P}^n \times T$  with composite map  $f : X \rightarrow T$  flat, meaning that for each  $x \in X$  with  $t = f(x)$ , the map  $\mathcal{O}_{T,t} \rightarrow f_*\mathcal{O}_{X,x}$  is a flat homomorphism of local rings. The following shows that flatness is the right notion:

**Theorem 2.1.** *If  $T$  is connected and  $X \subset \mathbb{P}^n \times T$ , then  $X \rightarrow T$  is flat if and only if the fibers  $X_t$  have constant Hilbert polynomial.*

**Example 2.2.** If  $C \subset \mathbb{P}^n$  is a curve and  $V \subset H^0(C, L)$  gives the linear system  $\mathbb{P}V$ , then the total family  $D \subset C \times \mathbb{P}V \subset \mathbb{P}^n \times \mathbb{P}V$  with fibers  $D_s = (s)_0$  is a flat family.

The amazing thing is that there is a scheme that captures all such flat families at once. Grothendieck showed [1] that all flat families with fixed Hilbert polynomial  $p(t) \in \mathbb{Q}[t]$  are captured by one universal flat family in the following theorem.

**Theorem 2.3.** *There is a scheme  $\text{Hilb}_{p(m)}^n$  and flat family (1)  $X \subset \mathbb{P}^n \times \text{Hilb}_{p(m)}^n$  such that for each flat family (2)  $C \subset \mathbb{P}^n \times T \rightarrow T$  with fibers having Hilbert polynomial  $p(t)$ , (2) arises from (1) as a pullback from a unique map  $T \rightarrow \text{Hilb}_{p(m)}^n$ .*

**Remark 2.4.** The Hilbert schemes  $\text{Hilb}_{p(t)}^n$  have been both objects of study and a useful tool since their arrival in 1960. A few comments about their nature:

- (a) Grothendieck proved that  $\text{Hilb}_{p(t)}^n$  is projective over  $\text{Spec } \mathbb{Z}$ , so every flat family is encoded by a projective scheme with integer coefficients.
- (b) Over a field  $k$ , Hartshorne showed [3] that  $\text{Hilb}_{p(t)}^n$  is connected for each  $p(t)$ , but examples show it can have many irreducible components.
- (c) The universal property implies that  $\text{Hilb}_{p(t)}^n$  is unique up to unique isomorphism.
- (d) Taking  $T = \text{Spec } k$  shows that there is a bijection

$$\{\text{Points of } \text{Hilb}_{p(t)}^n\} \leftrightarrow \{\text{Closed subschemes } X \subset \mathbb{P}^n \text{ with Hilbert polynomial } p(t)\}$$

**Exercise 2.5.** Prove that  $\text{Hilb}_1^n = \mathbb{P}^n$  with universal family  $X = \Delta(\mathbb{P}^n) \subset \mathbb{P}^n \times \mathbb{P}^n$ .

**Notation 2.6.** For degree  $d$  and genus  $g$ , we write  $\text{Hilb}_{d,g}^n = \text{Hilb}_{dm+1-g}^n$ .

**Example 2.7.** The twisted cubic curve  $C \subset \mathbb{P}^3$  has ideal  $I = (y^2 - xz, xw - yz, yw - z^2)$  generated by the maximal minors of  $M = \begin{pmatrix} z & y & x \\ w & z & y \end{pmatrix}$ . The Hilbert-Burch theorem gives a free resolution  $0 \rightarrow S(-3)^2 \xrightarrow{M} S(-2)^3 \rightarrow I_C \rightarrow 0$ . We vary entries of  $M$  to give flat families of curves parametrized by  $t \in \mathbb{A}^1$  and evaluate the limiting curve as  $t \rightarrow 0$ .

- (a)  $M_t = \begin{pmatrix} z & y & x \\ w & zt & y \end{pmatrix} \Rightarrow I_{C_0} = (wy, xw - yz, y^2)$ , so  $C_0$  is the union of  $x = y = 0$  and doubling of  $w = y = 0$  on  $Q$ .
- (b)  $M_t = \begin{pmatrix} z & y & xt \\ w & z & y \end{pmatrix} \Rightarrow I_{C_0} = (wy - z^2, zy, y^2)$ , a triple line on the quadric cone.
- (c)  $M_t = \begin{pmatrix} z & y & xt \\ wt & z & y \end{pmatrix} \Rightarrow I_{C_0} = (z^2, zy, y^2) = (x, y)^2$ , the thick triple line.
- (d) Projection from a point gives a nodal plane cubic with a spatial embedded point at the node [4, III, Ex. 9.8.4] (see also [7]).

**Example 2.8.** Work of Piene and Schlessinger [7] describes  $\text{Hilb}_{3,0}^3$ . It has two irreducible components: (a) the 12-dimensional closure of the family of twisted cubic curves and (b) the 15-dimensional family whose general member is the union of a plane cubic and an isolated point. They meet transversely in an irreducible subvariety of dimension 11.

### 3. THE MODULI SPACE $M_g$ OF CURVES OF GENUS $g \geq 2$

We construct a scheme  $M_g$  whose closed points are in bijective correspondence with isomorphism classes of curves of genus  $g$ .

**Example 3.1.** If  $g = 0$ , then applying Corollary 1.13 to  $D = P \in C$  gives a closed immersion  $C \rightarrow \mathbb{P}^1$  which must be an isomorphism, so  $M_0$  is a point.

**Example 3.2.** The  $j$ -invariant shows that  $M_1 = \mathbb{A}^1$ .

In view of Examples 3.1 and 3.2 we assume  $g \geq 2$ . We need a way to produce all the curves and recognize isomorphism classes. This is provided by the following:

**Observations 3.3.** Let  $C$  be a curve of genus  $g \geq 2$ .

- (a) Fix  $n \geq 3$ . Then the  $n$ -canonical complete linear system  $|nK_C|$  gives a closed embedding  $C \hookrightarrow \mathbb{P}^N$  by Corollary 1.13, with  $N = n(2g-2) - g = (2n-1)(g-1) - 1$ . The image is an  $n$ -canonical curve of degree  $d = n(2g-2)$  and genus  $g$ .
- (b) If  $j : C \hookrightarrow \mathbb{P}^N$  is an  $n$ -canonical embedding and  $\Psi \in \text{Aut } \mathbb{P}^N$ , then  $\Psi \circ j$  is also an  $n$ -canonical embedding.
- (c) If  $\varphi : C' \xrightarrow{\sim} C$  is an isomorphism, then the composite  $C' \rightarrow C \rightarrow \mathbb{P}^N$  is also an  $n$ -canonical embedding because  $\varphi^*K_C = K_{C'}$ . It follows that each isomorphism class of genus  $g$  curves appears a single  $\text{Aut } \mathbb{P}^N$ -orbit.

Taking  $G = \text{Aut } \mathbb{P}^N = \text{PGL}(N+1)$ , this leads to the following.

**Construction 3.4.** Let  $U \subset \text{Hilb}_{n(2g-2),g}^N$  be the open subset of smooth connected curves and  $K \subset U$  the subset of  $n$ -canonical curves. Then  $G$  acts on  $K$  and the set  $K/G$  of  $G$ -orbits is in bijective correspondence with isomorphism classes of genus  $g$  curves.

**3.1. Geometric Invariant theory.** Given a group  $G$  acting on a variety  $X$ , geometric invariant theory [5] establishes conditions under which the orbit space  $X/G$  exists as the points of an algebraic variety. Here are a summary of some results from this area. Below we take  $G$  to be a *reductive* group, of which  $G = \text{PGL}(N+1)$  is an example.

**3.2. Affine case.** If  $X = \text{Spec } A$  is affine, things work out well: the action of  $G$  on  $X$  induces an action  $G \times A \rightarrow A$ , when we can define  $A^G = \{a \in A : ga = a \forall g \in G\}$  and the map  $\phi : \text{Spec } A \rightarrow Y = \text{Spec } A^G$  does the job. In particular, for any open  $U \subset Y$  for which the  $G$ -orbits in  $\phi^{-1}(U)$  are closed, then  $U = \phi^{-1}(U)/G$  is the space of orbits.

**3.3. Projective case.** If  $X = \text{Proj } R$  where  $R$  is graded, things are more difficult.

3.3.1. *GIT Semistability.* Assuming that  $G$  acts on  $X$  *linearly*, meaning through the action of the general linear group on  $R_1$ , one can again form the  $G$ -invariant subring  $R^G \subset R$  and the map  $\pi : X \rightarrow Y = \text{Proj } R^G = X//G$ , but typically  $\pi$  will not be defined for all  $x \in X$ , so we define the semistable locus  $X^{ss} \subset X$  to be the open set where it is defined. Concretely  $x \in X$  is semistable if there is some  $f \in R^G$  with  $x \in X_f$ , otherwise  $x$  is nonstable. Then we get a morphism  $X^{ss} \rightarrow X//G$ .

3.3.2. *GIT stability.* Unfortunately when we use semistable points as above, the morphism might collapse distinct orbits together in  $X//G$ . To remedy this situation, we define  $x \in X^{ss}$  to be stable if each orbit  $Gx$  is closed in  $X^{ss}$  and  $\dim Gx = \dim G$  (in other words,  $|\text{Stab}(x)| < \infty$ ). If  $X^s \subset X^{ss}$  is the subset of stable points, then  $\pi : X^s \rightarrow X//G$  separates the orbits and  $X^s \rightarrow \pi(X^s)$  is the orbit space we are seeking.

The condition  $|\text{Stab}(x)| < \infty$  says that  $|\text{Aut } C| < \infty$  for the corresponding curve  $C$ . For a curve  $C$  of genus  $g$ , note the following:

- (1)  $g \geq 2 \Rightarrow |\text{Aut } C| < \infty$  [4, IV, Exercise 2.5].
- (2) If  $g = 1$ , only finitely many automorphisms fix  $P \in C$  [4, IV, Proposition 4.7].
- (3) If  $g = 0$ , automorphisms are determined by three points [4, I, Exercise 6.6].

With this in mind, we describe two types of nodal curves that also have finite automorphism groups, these are the Deligne-Mumford-stable curves:

- (a) the curves  $C$  such that every smooth rational component meets the other components in at least 3 points.
- (b) the curve  $C$  such that every rational component of the normalization of  $C$  has at least 3 points lying over the singular points of  $C$ .

Finally we can state the main result. We have defined  $K \subset U \subset \text{Hilb}$  above. Enlarge  $K$  to  $\tilde{K}$  by adding the curves of types (a) and (b) shown above and let  $\tilde{K}^{ss} = \tilde{K} \cap \text{Hilb}^{ss}$ .

**Theorem 3.5.** *Assume  $n \geq 5$  in the construction above.*

- (1)  $\tilde{K}^{ss}$  is a smooth closed subscheme of  $\text{Hilb}$ .
- (2)  $\tilde{K}^{ss} = \tilde{K}^s$ : these curves are already GIT stable.
- (3)  $\tilde{K}^{ss} = \tilde{K}$ .

*In particular,  $\tilde{K}^{ss}/G$  is a projective variety containing  $K/G$  as a dense open subset.*

## REFERENCES

- [1] A. Grothendieck, Techniques de construction et théorèmes d'existence en géométrie algébrique, IV: Les schémas de Hilbert, in Sem. Bourbaki, 1960-1961, Exposé 221, Benjamin, 1966.
- [2] J. Harris and I. Morrison, *Moduli of Curves*, GTM **187**, Springer-Verlag, 1998.
- [3] R. Hartshorne, Connectedness of the Hilbert scheme, Inst. Hautes Études Sci. Publ. Math. 29 (1966) 5–48.
- [4] R. Hartshorne, *Algebraic Geometry*, GTM **52**, Springer-Verlag, 1977.
- [5] J. Fogarty, F. Kirwan and D. Mumford, *Geometric invariant theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete 34, Folge 3, third edition, Springer-Verlag, 1994.
- [6] P. Newstead, *Introduction to moduli problems and orbit spaces*, Tata Institute of Fundamental Research Lectures on Mathematics and Physics, 51, Tata Institute of Fundamental Research, Bombay; by the Narosa Publishing House, New Delhi, 1978.

- [7] R. Piene and Schlessinger, On the Hilbert scheme compactification of the space of twisted cubics, Amer. J. Math. 107 (1985) 761–774.