

NOTICE: This paper, *Intersection homology with field coefficients: K-Witt spaces and K-Witt bordism*, contains an error in the characteristic 2 case. A copy of the Corrigendum and follow-up work in *K-Witt bordism in characteristic 2* have been appended at the end of the paper.

Greg Friedman

Intersection homology with field coefficients: K -Witt spaces and K -Witt bordism

Greg Friedman
Texas Christian University

July 23, 2008

Abstract

We construct geometric examples of pseudomanifolds that satisfy the Witt condition for intersection homology Poincaré duality with respect to certain fields but not others. We then compute the bordism theory of K -Witt spaces for an arbitrary field K , extending results of Siegel for $K = \mathbb{Q}$.

2000 Mathematics Subject Classification: 55N33, 57Q20, 57N80

Keywords: intersection homology, universal coefficient theorem, Witt space, Witt bordism

Contents

1	Introduction	2
2	Background material	4
3	Oddities of finite coefficients	7
3.1	Violations of the universal coefficient theorem	8
3.2	K -Witt spaces that are not K' -Witt spaces	10
4	K-Witt bordism groups	15
4.1	Preliminaries	16
4.2	Reduction to prime fields	16
4.3	Witt bordism over \mathbb{Z}_p	18
4.4	Witt bordism over finite fields	20
4.5	\mathbb{Z}_p -Witt bordism as a generalized homology theory	21

1 Introduction

This paper consists of two related parts: In the first part, we provide some examples of the phenomena that arise when considering intersection homology over coefficient groups with torsion, including various forms of violation of the universal coefficient theorem and spaces that satisfy the *Witt condition* for certain fields but not others (and hence possess Poincaré duality with respect to certain fields but not others). In the second part, we compute the bordism theory of spaces that are K -Witt for an arbitrary field K . Let us explain what all this means by providing some rough definitions and context; more precise background details can be found below in Section 2.

Intersection homology groups were developed by Goresky and MacPherson in [13] for the purpose of extending Poincaré duality and ensuing invariants to non-manifold spaces. In [13], this is accomplished, *over rational coefficients*, for the class of (compact, oriented) piecewise-linear pseudomanifolds, a class of spaces including all projective complex varieties. Using sheaf theory, this duality was later expanded [14] to the broader class of topological pseudomanifolds and to coefficients in any field. Other generalizations followed; see, e.g. [7].

The dual pairing of intersection homology, in generality, pairs intersection homology groups with dual indices (as in the familiar case of manifolds) *and* with dual sets of perversity parameters. Thus the general duality result for an n -dimensional compact oriented pseudomanifold asserts that there is a perfect (nonsingular) pairing

$$I^{\bar{p}}H_i(X; K) \times I^{\bar{q}}H_{n-i}(X; K) \rightarrow K.$$

Here IH denotes the intersection homology groups, K is a coefficient field, and \bar{p} and \bar{q} are dual sets of *perversity parameters* that occur as one of the inputs to the theory; see Section 2, below, for details.

Ideally, however, one would like a little more. For $2k$ -dimensional spaces, one would like a $(-1)^k$ -symmetric self-pairing on $I^{\bar{p}}H_k(X; K)$. This would yield signatures, elements of Witt groups, and other further algebraic information. Unfortunately, this is not possible in general because there are no self-dual perversities, but there are dual “middle perversities,” \bar{m} and \bar{n} , and certain spaces such that $I^{\bar{m}}H_i(X; K) \cong I^{\bar{n}}H_i(X; K)$, in which case we obtain the desired middle dimensional form.

It was recognized early on (right in [13]) that pseudomanifolds with only even codimension singularities possess this form of self-duality. This class was soon generalized by Siegel [27] to a class of spaces he dubbed Witt spaces, and which we will more specifically call \mathbb{Q} -Witt spaces. These spaces are identified by certain local intersection homology conditions, and they possess the middle-dimensional self-duality over \mathbb{Q} . Siegel further computed the bordism groups of these spaces, showing that in nontrivial cases they equal the Witt group $W(\mathbb{Q})$ - hence the name “Witt spaces” - and that the resulting bordism homology theory provides a geometric formulation of KO -homology at odd primes. Further work in this direction was done by Pardon [25], who computed the bordism groups of the more restrictive integer “Poincaré Duality spaces” of Goresky and Siegel [15] and related these groups to computations of $[X, G/TOP]$.

Banagl [2] has since extended duality even further by identifying conditions on which non-Witt spaces possess self-duality (conditions equivalent to the existence of certain towers of Lagrangian structures on strata), but Witt spaces remain an important class of examples defined by a relatively tractable condition.

This brings us to intersection homology with coefficients. Unlike ordinary homology theories, intersection homology does not, in general, possess a universal coefficient theorem (though Goresky and Siegel [15] have shown, using the Deligne sheaf formulation of intersection homology, that a universal coefficient sequence will occur if a space possesses certain local torsion properties). Thus, intersection homology groups with different coefficients must be treated in their own right, without any clear connections between them. In particular, spaces that satisfy intersection homology Poincaré duality with one set of coefficients may not possess duality with other sets of coefficients.

Our first goal is to provide some examples of these phenomena. We produce concrete examples of spaces where the universal coefficient theorem breaks down (in different ways), and we present spaces that are K -Witt (and hence possess self-duality) with respect to some coefficient fields K but not others. Our arguments and constructions are purely geometric, avoiding sheaf theory in favor of hands-on examination of intersection chains.

The following facts will be demonstrated throughout Section 3.2 (except for the first, which is shown in Section 4.2):

Theorem 1.1. *Let K denote a field, and let \mathbb{Z}_p denote the field of p elements.*

1. *If K has characteristic $p > 0$, then X is K -Witt if and only if X is \mathbb{Z}_p -Witt; if K has characteristic 0, then X is K -Witt if and only if X is \mathbb{Q} -Witt.*
2. *If $n > 4$ and P is a finite set of primes, then there is a compact orientable n -dimensional pseudomanifold that is \mathbb{Z}_p -Witt for any $p \in P$ but that is not \mathbb{Q} -Witt and not \mathbb{Z}_p -Witt for $p \notin P$.*
3. *If $n > 4$ and P is a finite set of primes, then there are \mathbb{Q} -Witt spaces that are not \mathbb{Z}_p -Witt for any $p \in P$ and are \mathbb{Z}_p -Witt for $p \notin P$.*
4. *If X is a 3- or 4-dimensional \mathbb{Z}_p -Witt space, then X is a \mathbb{Q} -Witt space.*
5. *If X is a 3- or 4-dimensional \mathbb{Q} -Witt space, then X is a \mathbb{Z}_p -Witt space for any $p \neq 2$. If X is also \mathbb{Q} -orientable, then it is also a \mathbb{Z}_2 -Witt space. However, there are non-orientable 3- and 4-dimensional PL \mathbb{Q} -Witt spaces that are not \mathbb{Z}_2 -Witt spaces.*
6. *All 0-, 1-, and 2-dimensional pseudomanifolds are K -Witt for all K .*

We also find examples of $4k$ -dimensional pseudomanifolds demonstrating conditions (2) and (3) that not just satisfy or fail to satisfy the appropriate Witt conditions but that also definitively possess or fail to possess the associated dualities in nontrivial ways.

Finally, in the second half of the paper, Section 4, we follow Siegel [27] by computing the bordism groups $\Omega_n^{K\text{-Witt}}$ of oriented K -Witt spaces for any coefficient field K as well as identifying the resulting generalized homology theories. We show the following theorems

(bear in mind item (1) of the preceding theorem, which implies that $\Omega_n^{K-\text{Witt}} \cong \Omega_n^{K'-\text{Witt}}$ if K and K' have the same characteristic):

Theorem 1.2 (Theorem 4.5). *Let $p \neq 0$ be a prime, and let $W(\mathbb{Z}_p)$ denote the Witt group of symmetric bilinear forms over the field \mathbb{Z}_p of p elements. Then*

1. $\Omega_0^{\mathbb{Z}_p-\text{Witt}} \cong \mathbb{Z}$,
2. for $n \not\equiv 0 \pmod{4}$, $\Omega_n^{\mathbb{Z}_p-\text{Witt}} = 0$, and
3. for $n \equiv 0 \pmod{4}$, $n > 0$, the homomorphism $w : \Omega_n^{\mathbb{Z}_p-\text{Witt}} \rightarrow W(\mathbb{Z}_p)$ that assigns to X^n the intersection form on $I^{\bar{n}}H_{n/2}(X; \mathbb{Z}_p)$ is an isomorphism.

Theorem 1.3 (Theorem 4.10). *As a homology theory on CW complexes, \mathbb{Z}_p -Witt bordism splits as a direct sum of ordinary homology with coefficients. In particular,*

$$\Omega_n^{\mathbb{Z}_p-\text{Witt}}(X) \cong \bigoplus_{r+s=n} H_r(X; \Omega_s^{\mathbb{Z}_p-\text{Witt}}).$$

Acknowledgment. I thank Shmuel Weinberger for suggesting that it would be interesting to study intersection homology with coefficients, in general, and for asking about extending Siegel's results to \mathbb{Z}_p -Witt bordism, in particular. In the course of chasing down Theorem 4.10, I had the pleasure of several fruitful correspondences and therefore owe great thanks to John McCleary, Jim McClure, Andrew Ranicki, Yuli Rudyak, Larry Taylor, and Shmuel Weinberger.

2 Background material

In this section we provide the relevant background for the rest of the paper.

Pseudomanifolds. We will work entirely in the class of piecewise linear (PL) spaces, although intersection homology can be defined more broadly on topological pseudomanifolds (see [14]).

We recall (see [13]) that a PL stratified space X is a PL space equipped with a filtration (compatible with the PL structure)

$$X = X^n \supset X^{n-1} \supset X^{n-2} \supset \dots \supset X^0 \supset X^{-1} = \emptyset$$

such that for each point $x \in X_i = X^i - X^{i-1}$, there exists a lower-dimensional compact PL stratified pseudomanifold L , a compatible filtration of L

$$L = L^{n-i-1} \supset \dots \supset L^0 \supset L^{-1} = \emptyset,$$

and a *distinguished neighborhood* U of x such that there is a PL homeomorphism

$$\phi : \mathbb{R}^i \times c(L) \rightarrow U$$

that takes $\mathbb{R}^i \times c(L^{j-1})$ onto $X^{i+j} \cap U$. Here cL denotes the open cone on L . In other words, each point has a neighborhood that is a trivial bundle of cones on a lower-dimensional stratified space.

A PL stratified space X is a PL stratified pseudomanifold if $X^{n-1} = X^{n-2}$ and $X - X^{n-1}$ is dense in X . A space is called a PL pseudomanifold if it can be given the structure of a PL stratified pseudomanifold by some choice of stratification.

The subspace $X_i = X^i - X^{i-1}$ is called the i th *stratum*, and, in particular, it is a (possibly empty) PL i -manifold. L is called the *link* of the component of the stratum. Note that we do not allow a codimension 1 stratum. There are various technical reasons for this, including the avoidance of “pseudomanifolds with boundary” (see, e.g., [11, 10], where this issue is treated in detail); however, we will revisit this idea in our discussion of bordism in Section 4. $X^{n-1} = X^{n-2}$ is often referred to as the *singular locus* and denoted Σ . A PL stratified pseudomanifold X is *oriented* if $X - \Sigma$ is oriented as a manifold.

Intersection homology. Intersection homology, due to Goresky and MacPherson [13], is a topological invariant of pseudomanifolds (in particular, it is invariant under choice of PL structure or stratification - see [14], [5], [17]). It possesses a definition via sheaf theory, which is important (indeed crucial) for many applications, but the original definition was given as the homology of a subcomplex of the complex $C_*(X)$ of PL chains on X . This $C_*(X)$ is a direct limit $\varinjlim_{T \in \mathcal{T}} C_*^T(X)$, where $C_*^T(X)$ is the simplicial chain complex with respect to the triangulation T , and the direct limit is taken with respect to subdivision within a family of triangulations compatible with each other under subdivision and compatible with the filtration of X . In fact, while it is convenient to work with these PL chains, one can also work with simplicial chains, supposing a fine enough triangulation of X (see the appendix to [20]).

Intersection chain complexes are subcomplexes of $C_*(X)$ defined with regard to *perversity parameters* $\bar{p} : \mathbb{Z}^{\geq 2} \rightarrow \mathbb{Z}$ that are required to satisfy $\bar{p}(2) = 0$ and $\bar{p}(k) \leq \bar{p}(k+1) \leq \bar{p}(k) + 1$. We think of the perversity as taking the codimensions of the strata of X as input. The output tells us the extent to which chains in the intersection chain complex will be allowed to intersect that stratum. Thus a simplex σ in $C_i(X)$ (represented by a simplex in some triangulation) is deemed \bar{p} -allowable if $\dim(\sigma \cap X^{n-k}) \leq i - k + \bar{p}(k)$, and a chain $\xi \in C_i(X)$ is \bar{p} -allowable if every simplex with non-zero coefficient in ξ or $\partial\xi$ is allowable as a simplex. The allowable chains constitute the chain complex $I^{\bar{p}}C_*(X)$, and the \bar{p} -perversity intersection homology groups $I^{\bar{p}}H_*(X)$ are the homology groups of this chain complex. Note that if M is a manifold, then $I^{\bar{p}}H_*(M) \cong H_*(M)$. This is not obvious if M is stratified in an interesting way, but it follows from the topological invariance of intersection homology groups, which implies that $I^{\bar{p}}H_*(M)$ may be computed from the trivial filtration $M \supset \emptyset$.

For more general background on intersection homology, we urge the reader to consult the expositions by Borel, et. al. [5] or Banagl [3]. For both background and application of intersection homology in various fields of mathematics, the reader should see Kirwan and Woolf [18].

Intersection homology with coefficients. The definition of intersection homology with coefficients is given analogously so that $I^{\bar{p}}C_*(X; G)$ is the subcomplex of $C_*(X; G) \cong C_*(X) \otimes G$, again consisting of chains ξ such that every simplex with non-zero coefficient in ξ or $\partial\xi$ is allowable as a simplex. However, a critical point to observe is that, in general, $I^{\bar{p}}C_*(X; G)$ is not isomorphic to $I^{\bar{p}}C_*(X) \otimes G$. It is true that a simplex with nonzero coefficient in a chain $\xi \in I^{\bar{p}}C_*(X; G)$ is allowable or not depending only on the simplex itself and not the coefficient. However, which simplices appear with non-zero coefficient in $\partial\xi$ might depend strongly on the coefficients being used.

For example, consider a chain of the form $\xi = \sum_i \sigma_i$ over some collection of oriented simplices, each with coefficient 1. The allowability of each of the simplices σ_i is independent of whether we think of ξ as a chain in $C_i(X)$ or $C_i(X; \mathbb{Z}_2)$. Now suppose each σ_i in ξ is allowable, and consider $\partial\xi$. Suppose that $\partial\xi = 2\eta$ for some chain η . It is possible in $C_i(X)$ that η may contain simplices that are not allowable. However, in $C_i(X; \mathbb{Z}_2)$, $\partial\xi = 0$, and the allowability conditions are satisfied vacuously.

Thus when working with coefficients, the obvious homomorphism $I^{\bar{p}}C_*(X) \otimes G \rightarrow I^{\bar{p}}C_*(X; G)$ is an injection, but it is not, in general, a surjection. These considerations, of course, have the potential to affect the intersection homology groups quite radically. For example, the universal coefficient theorem is not generally valid for intersection homology. In the next section, we turn to concrete examples that demonstrate geometrically what can go wrong.

The cone formula. Perhaps the most important concrete computation in intersection homology is the formula for the intersection homology of an open cone. If L is an n -dimensional compact pseudomanifold, then the open cone cL is stratified so that $(cL)^0$ is the cone vertex and, for $i > 0$, $(cL)^i = L^{i-1} \times (0, 1) \subset cL$. Then the intersection homology of the cone cL is given as follows:

$$I^{\bar{p}}H_i(cL; G) \cong \begin{cases} 0, & i \geq n - \bar{p}(n + 1), \\ I^{\bar{p}}H_i(L; G), & i < n - \bar{p}(n + 1). \end{cases}$$

This formula comes from direct consideration of the definition of the intersection chain complex and the fact that the dimension of the intersection of a simplex with the cone vertex can be at most 0. See [5, Section 1] for more details.

It is also useful to have the formula for the intersection homology of a suspension, which comes from the cone formula and a Mayer-Vietoris argument (see¹ [17]). If X is a compact n -dimensional pseudomanifold with suspension SX , then

$$I^{\bar{p}}H_i(SX; G) \cong \begin{cases} I^{\bar{p}}H_{i-1}(X; G), & i > n - \bar{p}(n + 1), \\ 0, & i = n - \bar{p}(n + 1), \\ I^{\bar{p}}H_i(X; G), & i < n - \bar{p}(n + 1). \end{cases}$$

¹The formula presented here is slightly simpler than the one in [17] since we allow only traditional perversities, not “loose” perversities.

Witt spaces. The chief interest (at least originally) in intersection homology is that, with field coefficients, it satisfies Poincaré Duality. More specifically, assume that X^n is a compact, oriented, and *irreducible* (meaning $X - \Sigma$ is connected) PL pseudomanifold, and let F be a field and \bar{p} and \bar{q} dual perversities (meaning that $\bar{p}(k) + \bar{q}(k) = k - 2$ for all k). Then there is a nondegenerate pairing $I^{\bar{p}}H_i(X; F) \otimes I^{\bar{q}}H_{n-i}(X; F) \rightarrow F$, defined via the intersection pairing on intersection chains in general position. We refer the reader to [13, 14, 5, 8] for more details. While this is good, one would like something even better, a condition that guarantees a self-pairing between middle-dimensional intersection homology groups for even-dimensional manifolds. This is what the Witt spaces provide.

Let \bar{m} and \bar{n} be the lower and upper middle perversities given by $(0, 0, 1, 1, 2, 2, \dots)$ and $(0, 1, 1, 2, 2, 3, \dots)$, i.e. $\bar{m}(k) = \lfloor \frac{k-2}{2} \rfloor$ and $\bar{n}(k) = \lfloor \frac{k-1}{2} \rfloor$. These are dual perversities, and it is not hard to check from the definitions that if X^{2n} is compact and oriented and has nonempty strata only of even dimension, then $I^{\bar{m}}H_*(X; F) \cong I^{\bar{n}}H_*(X; F)$. So in this case there is a $(-1)^n$ -symmetric form $I^{\bar{m}}H_n(X; F) \otimes I^{\bar{m}}H_n(X; F) \rightarrow F$. When X has dimension $4n$ this yields signatures, etc; see [13]. A weaker condition on X^n that yields the same outcome is the *F-Witt condition*, which assumes that $I^{\bar{m}}H_k(L; F) = 0$ for each link L^{2k} of each stratum of dimension $n - 2k - 1$, $k > 0$. In this case, it follows once again that $I^{\bar{m}}H_*(X; F) \cong I^{\bar{n}}H_*(X; F)$ (see [5]), and we obtain middle dimensional pairings [14]. The Witt condition turns out to be independent of the choice of stratification of X [27] and so it is a property of the pseudomanifold X , itself.

In keeping with the conventions of [13] and [27], we will call an *oriented compact irreducible PL stratified pseudomanifold* satisfying the *F-Witt condition* an *F-Witt space*. The orientation condition is implicit in [27] based on the definition of pseudomanifold given in [13]. If we need to refer to a nonorientable pseudomanifold satisfying the Witt condition, we will call it explicitly a “non-orientable Witt space.”

Witt spaces were so-named by Siegel due to his observation in [27] that the bordism groups of $4k$ -dimensional Witt spaces, $k > 0$, are isomorphic to the Witt group $W(\mathbb{Q})$. We recall that, for any commutative ring with unity R , the Witt group $W(R)$ is the group of equivalence classes of symmetric inner product spaces over R under the equivalence relation $X \sim X'$ if $X \oplus S \cong X' \oplus S'$, where S and S' are split inner product spaces, i.e. S and S' possess direct summands N and N' that are equal to their own orthogonal complements. The group operation is orthogonal direct sum. $W(R)$ can also be given a ring structure via the tensor product. For more on Witt groups, see [22, 19].

3 Oddities of finite coefficients

In this section, we begin with some simple examples of the violation of the universal coefficient theorem for intersection homology. We move on to more complex examples that are then used to construct spaces that satisfy Witt conditions with respect to certain fields but not others.

3.1 Violations of the universal coefficient theorem

In [15], Goresky and Siegel used sheaf machinery to prove that \bar{p} -perversity intersection homology satisfies the universal coefficient theorem for an abelian group G on a pseudomanifold X if X is *locally \bar{p} -torsion free*. This condition means that if L is the link of a stratum of X of codimension c , then the abelian group $I^{\bar{p}}H_{c-2-\bar{p}(c)}(L)$ is torsion free. While the proof of the theorem in [15] involves the axiomatic sheaf formulation of intersection homology, one can work directly with chains to find examples of the trouble that can arise if this torsion condition is violated. In this section, we provide several such examples of varying degrees of complexity.

A simple example of violation of universal coefficients. As a first example of the violation of the universal coefficient theorem in intersection homology, consider $X = c(\mathbb{R}P^2)$, the open cone on $\mathbb{R}P^2$, and suppose that $\bar{p}(3) = 0$. The link of the singular vertex v of X is $L = \mathbb{R}P^2$, and $I^{\bar{p}}H_1(L) = H_1(L) = \mathbb{Z}_2$. So X is not locally \bar{p} -torsion free.

We compute from the cone formula (see Section 2):

$$I^{\bar{p}}H_i(c\mathbb{R}P^2) \cong \begin{cases} 0, & i \geq 2 - \bar{p}(3) = 2, \\ I^{\bar{p}}H_i(\mathbb{R}P^2), & i < 2. \end{cases}$$

In particular, $I^{\bar{p}}H_1(X) \cong \mathbb{Z}_2$ and $I^{\bar{p}}H_2(X) \cong 0$.

Similarly, since the cone formula holds for any coefficients,

$$I^{\bar{p}}H_i(c\mathbb{R}P^2; \mathbb{Z}_2) \cong \begin{cases} 0, & i \geq 2 - \bar{p}(3) = 2, \\ I^{\bar{p}}H_i(\mathbb{R}P^2; \mathbb{Z}_2), & i < 2, \end{cases}$$

and so also $I^{\bar{p}}H_1(X; \mathbb{Z}_2) \cong \mathbb{Z}_2$ and $I^{\bar{p}}H_2(X; \mathbb{Z}_2) \cong 0$.

But this violates the universal coefficient theorem, which would predict that $I^{\bar{p}}H_2(X; \mathbb{Z}_2)$ would equal $(I^{\bar{p}}H_2(X) \otimes \mathbb{Z}_2) \oplus (I^{\bar{p}}H_1(X) * \mathbb{Z}_2) \cong \mathbb{Z}_2$.

We can see in this example a situation in which a chain that would not be allowable in integer intersection homology becomes allowable in intersection homology with coefficients – recall from Section 2 that it is this effect that is ultimately responsible for the violation of the universal coefficient theorem. Specifically, consider the standard cell decomposition² of $\mathbb{R}P^2$ with one cell in each dimension, and let x be the 1-cell that represents the generator of $H_1(\mathbb{R}P^2)$. Similarly, let y be the 2-cell with $\partial y = 2x$. y does not represent an integer homology class because it is not a cycle, but with coefficients in \mathbb{Z}_2 , $\partial y = 0$ and $[y]$ represents the nontrivial class in $H_2(\mathbb{R}P^2) \cong \mathbb{Z}_2$. This is precisely the term coming from the torsion product in the universal coefficient theorem in ordinary homology. Now, in $c\mathbb{R}P^2$, consider the 3-chain cy determined by the cone on y . This is *not* an allowable chain with integer coefficients because even though $\dim(cy) \cap X^0 = 0 \leq 3 - 3 + \bar{p}(3) = 0$, we have $\partial(cy) = y - c\partial y = y - 2x$, and this 2-chain intersects the cone point, which is not allowed with this perversity. However, with \mathbb{Z}_2 coefficients, $\partial(cy) = y$, which does not intersect the

²We give a description using cells, but this argument could also be made simplicially.

cone point; thus cy is allowable and kills the cycle y . It is interesting to note that with \mathbb{Z} coefficients y is not even a cycle to begin with!

A “worse” violation of universal coefficients. In the last example, we saw that intersection homology can violate the universal coefficient theorem when the Goresky-Siegel local torsion condition is violated. More specifically, the expected torsion product summands did not materialize. In the next example, we see that something more unexpected can happen: the tensor product terms might also vanish. In particular, we will construct spaces that have non-trivial integer intersection homology in their middle dimensions but whose intersection homology with finite coefficients vanishes in the same dimensions.

These examples were motivated initially by applying a construction that Siegel uses with rational coefficients in [27]. We also follow the arguments of Haefliger from [5, Section I.5.3] for computing the intersection homology of a Thom space.

Consider p copies of $\mathbb{C}P^2$ labeled $\mathbb{C}P_i^2$, $i = 1, \dots, p$. In each $\mathbb{C}P_i^2$, let V_i denote an embedded S^2 representing the generator of $H_2(\mathbb{C}P_i^2)$. Let $X = \#_{i=1}^p \mathbb{C}P_i^2$. We may assume that the V_i are disjoint within X , and we may form the connected sum $V = \#_{i=1}^p V_i \cong S^2$ embedded in X . The homology class $[V] \in H_2(\mathbb{C}P^2)$ is equal to $\sum [V_i]$. In particular, since the intersection number $[V_i] \cdot [V_i] = 1$ in $\mathbb{C}P_i^2$, we have $[V] \cdot [V] = p$.

Now let U denote a tubular neighborhood of V in X , and let \hat{U} denote the one point compactification of U . This is none other than the Thom space of the normal bundle to V in X . As shown in [5, Section I.5.3], the intersection homology of a Thom space is easy to compute. In general, if M is a compact n -manifold with boundary and $Y = M \cup_{\partial M} c(\partial M)$, a short calculation with the Mayer-Vietoris sequence and the cone formula demonstrates that

$$I^{\bar{p}} H_i(Y) \cong \begin{cases} H_i(Y), & i > n - \bar{p}(n) - 1, \\ \text{Im: } H_i(M) \rightarrow H_i(Y), & i = n - \bar{p}(n) - 1, \\ H_i(M), & i < n - \bar{p}(n) - 1. \end{cases}$$

Roughly speaking, in analogy with the cone formula, chains below a certain dimension are not allowed to intersect the distinguished cone point v , and in these dimensions the intersection homology is $H_*(Y - v) \cong H_*(M)$. In high dimensions, any chain is allowed, and the intersection homology is $H_*(Y) \cong H_*(M, \partial M)$. In the transition dimension, cycles cannot intersect v , but chains one dimension up can, and so we get the groups $\text{Im: } (H_i(M) \rightarrow H_i(Y)) \cong \text{Im: } (H_i(M) \rightarrow H_i(M, \partial M))$.

If now M is an r -disk bundle over a compact m -dimensional manifold B , then Y is the associated Thom space, and then we know $H_i(Y) \cong H_i(M, M - B) \cong H_{i-r}(B)$ by the Thom isomorphism theorem. In particular, $\text{Im}(H_i(M) \rightarrow H_i(Y)) \cong \text{Im}(H_i(B) \rightarrow H_i(Y)) \cong \text{Im}(H_i(B) \xrightarrow{e \cap} H_{i-r}(B))$, where e is the euler class of the bundle (see, e.g. [6, Section VI.12]).

In our case at hand, and using the lower middle perversity \bar{m} (see Section 2), we therefore have

$$I^{\bar{m}} H_i(\hat{U}) \cong \begin{cases} H_{i-2}(S^2), & i > 2, \\ \text{Im: } H_2(S^2) \xrightarrow{\cap e} H_0(S^2), & i = 2, \\ H_i(S^2), & i < 2. \end{cases}$$

So the nonzero groups are $I^{\bar{m}}H_0(\hat{U}) = \mathbb{Z}$, $I^{\bar{m}}H_4(\hat{U}) = \mathbb{Z}$, and $I^{\bar{m}}H_2(\hat{U}) = p\mathbb{Z} \cong \mathbb{Z}$, since the self-intersection number $[V] \cdot [V] = p$ is equal to the euler number.

Letting p be a prime, these same calculations hold over the field \mathbb{Z}_p of p elements except in this case we see that $I^{\bar{m}}H_0(\hat{U}; \mathbb{Z}_p) = \mathbb{Z}_p$, $I^{\bar{m}}H_4(\hat{U}; \mathbb{Z}_p) = \mathbb{Z}_p$, and, $I^{\bar{m}}H_2(\hat{U}; \mathbb{Z}_p) = p\mathbb{Z}_p = 0$. In addition, for a prime $p' \neq p$, we get $I^{\bar{m}}H_0(\hat{U}; \mathbb{Z}_{p'}) = \mathbb{Z}_{p'}$, $I^{\bar{m}}H_4(\hat{U}; \mathbb{Z}_{p'}) = p\mathbb{Z}_{p'} \cong \mathbb{Z}_{p'}$, and $I^{\bar{m}}H_2(\hat{U}; \mathbb{Z}_{p'}) = \mathbb{Z}_{p'}$.

Remark 3.1. More generally, if m is a positive composite integer and we perform the above construction with m copies of $\mathbb{C}P^2$, then we will have $I^{\bar{m}}H_2(\hat{U}; \mathbb{Z}_p) = 0$ for each prime p such that $p \mid m$ but $I^{\bar{m}}H_2(\hat{U}; \mathbb{Z}_{p'}) = \mathbb{Z}_{p'}$ for each prime p' such that $p' \nmid m$.

Obviously there is nothing particularly special here about having found our bundle within a connected sum of $\mathbb{C}P^2$ s. In fact, we can perform the same intersection homology computations starting with any n -bundle over an n -manifold and with an appropriate Euler number. However, our example also illustrates a more general procedure adapted from [27] for finding spaces with trivial middle-dimensional \mathbb{Z}_p intersection homology; see Remark 4.8, below.

3.2 K -Witt spaces that are not K' -Witt spaces

In this section, we construct spaces that are Witt with respect to certain fields but not Witt with respect to others, collectively demonstrating the assertions of Theorem 1.1 of the Introduction, with the exception of item (1), which is proven in Section 4. Recall that item (1) states that whether or not a space is K -Witt depends only on the characteristic of K ; hence in this section we consider only the fields \mathbb{Q} and \mathbb{Z}_p , the finite field of p elements.

Low dimensions. We first dispense with some low-dimensional considerations, establishing items (4), (5), and (6) of Theorem 1.1. We observe immediately that all 0- and 1-dimensional pseudomanifolds are manifolds, and hence K -Witt for all fields K , while 2-dimensional pseudomanifolds that are not manifolds can have only codimension 2 singularities and so are also K -Witt for all K . For dimensions 3 and 4, we have the following propositions.

Proposition 3.2. *Let X be a 3- or 4-dimensional \mathbb{Z}_p -Witt space. Then X is a \mathbb{Q} -Witt space.*

Proof. The only nontrivial even-dimensional links L in X must be 2-dimensional compact pseudomanifolds. But each of these is the union of a finite number of compact surfaces S_1, \dots, S_r , joined along a finite number of points (see [4]). Since intersection homology is invariant under normalizations³ (see [13, Section 4]), $I^{\bar{p}}H_1(L; \mathbb{Z}_p) \cong H_1(\amalg_i S_i; \mathbb{Z}_p)$, and so the result follows from the universal coefficient theorem for ordinary homology. \square

We also have the following converse:

Proposition 3.3. *If X is a 3- or 4-dimensional \mathbb{Q} -Witt space, then X is a \mathbb{Z}_p Witt space for any $p \neq 2$. If X is also \mathbb{Q} -orientable, then it is also a \mathbb{Z}_2 -Witt space. However, there are non- \mathbb{Q} -orientable 3- and 4-dimensional \mathbb{Q} -Witt spaces that are not \mathbb{Z}_2 -Witt spaces.*

³A pseudomanifold is normal if its links are connected, and every pseudomanifold is an image of a finite-to-one cover by a normal manifold, its normalization.

Proof. The proposition is true for $p \neq 2$, since if X is a 3- or 4-dimensional \mathbb{Q} -Witt space, then the two-dimensional links must consist of S^2 s and $\mathbb{R}P^2$ s glued along points, and these links will then have trivial \mathbb{Z}_p intersection homology in degree one, as well.

For $p = 2$, if a \mathbb{Q} -Witt space has a link that does involve at least one $\mathbb{R}P^2$, the space will not be \mathbb{Z}_2 -Witt. However, we claim such a space will not be \mathbb{Q} -orientable either. To see this, note that any such pseudomanifold must have distinguished neighborhoods of the form cL or $\mathbb{R}^1 \times cL$ and for which there is a map $\mathbb{R}P^2 \rightarrow L$ that is injective off of finitely many points. Any embedded curve representing a generator of $\pi_1(\mathbb{R}P^2)$ can be homotoped by a small homotopy to an embedded curve γ in L whose neighborhood in the distinguished neighborhood is homeomorphic to the product of a Möbius band with \mathbb{R}^1 or \mathbb{R}^2 . Thus tracing around γ in X must reverse orientation.

For examples of non-orientable 3- and 4-dimensional PL \mathbb{Q} -Witt spaces that are not \mathbb{Z}_2 -Witt, we can take the suspension and double suspension of $\mathbb{R}P^2$. \square

Remark 3.4. We note that the non- \mathbb{Q} -orientable \mathbb{Q} -Witt non- \mathbb{Z}_2 -Witt spaces also technically violate Siegel's definition of a \mathbb{Q} -Witt space in [27], since it implicitly uses the original Goresky-MacPherson definition of a pseudomanifold from [13], and this definition includes an orientability condition. In any event, there will be no rational perfect pairing on middle intersection homology, and this is what we like Witt spaces for.

Spaces that are \mathbb{Z}_p -Witt but not \mathbb{Q} -Witt or $\mathbb{Z}_{p'}$ -Witt for $p' \neq p$. We turn to demonstrating item (2) of Theorem 1.1 by next constructing, for $n > 4$, an n -dimensional \mathbb{Z}_p -Witt space, p -prime, that is neither \mathbb{Q} -Witt nor $\mathbb{Z}_{p'}$ -Witt for any prime $p' \neq p$. From the preceding section, there exists a 4-dimensional pseudomanifold \hat{U} whose $I^{\hat{m}}H_2(\hat{U}; \mathbb{Z}_p)$ vanishes but such that neither $I^{\hat{m}}H_2(\hat{U}; \mathbb{Q})$ nor $I^{\hat{m}}H_2(\hat{U}; \mathbb{Z}_{p'})$ vanishes for prime $p' \neq p$. It follows immediately that the suspension $S\hat{U}$ is \mathbb{Z}_p -Witt but not \mathbb{Q} -Witt and not $\mathbb{Z}_{p'}$ -Witt for any $p' \neq p$. By taking products with manifolds, $M \times S\hat{U}$, we obtain compact \mathbb{Z}_p -Witt spaces that are not Witt for fields of any other characteristic in all dimensions ≥ 5 .

However, to make these examples a bit more robust, we would like to find some $4k$ -dimensional spaces with these Witt properties and for which we can see directly that there is a nontrivial nonsingular middle intersection pairing over \mathbb{Z}_p but not over \mathbb{Q} or $\mathbb{Z}_{p'}$ for $p' \neq p$. For this, we use a slightly more elaborate starting point.

Consider a bundle of 2-planes over the torus $T^2 \cong S^1 \times S^1$ and with euler number $e = p$. These can be found, for example, by providing T^2 with a complex structure and then forming the complex line bundle associated to a divisor $p[x]$ for $x \in T^2$; see, e.g. [16]. Let Y be the associated Thom space. Then, by our computations above, we have

$$I^{\bar{m}} H_i(Y; \mathbb{Q}) \cong \begin{cases} H_{i-2}(T^2; \mathbb{Q}), & i > 2, \\ \mathbb{Q} \cong \text{Im}: H_2(T^2; \mathbb{Q}) \xrightarrow{\cap^e} H_0(T^2; \mathbb{Q}), & i = 2, \\ H_i(T^2; \mathbb{Q}), & i < 2, \end{cases}$$

$$\cong \begin{cases} \mathbb{Q}, & i = 4, \\ \mathbb{Q} \oplus \mathbb{Q}, & i = 3, \\ \mathbb{Q}, & i = 2, \\ \mathbb{Q} \oplus \mathbb{Q}, & i = 1, \\ \mathbb{Q}, & i = 0, \end{cases}$$

and with coefficients in $\mathbb{Z}_{p'}$, we obtain the same results with each \mathbb{Q} replaced by $\mathbb{Z}_{p'}$. Meanwhile,

$$I^{\bar{m}} H_i(Y; \mathbb{Z}_p) \cong \begin{cases} H_{i-2}(T^2; \mathbb{Z}_p), & i > 2, \\ 0 \cong \text{Im}: H_2(T^2; \mathbb{Z}_p) \xrightarrow{\cap^e} H_0(T^2; \mathbb{Z}_p), & i = 2, \\ H_i(T^2; \mathbb{Z}_p), & i < 2, \end{cases}$$

$$\cong \begin{cases} \mathbb{Z}_p, & i = 4, \\ \mathbb{Z}_p \oplus \mathbb{Z}_p, & i = 3, \\ 0, & i = 2, \\ \mathbb{Z}_p \oplus \mathbb{Z}_p, & i = 1, \\ \mathbb{Z}_p, & i = 0. \end{cases}$$

If α, β denote cycles generating $H_1(S^1 \times S^1; \mathbb{Z}_p)$, then α and β also generate $I^{\bar{m}} H_1(Y; \mathbb{Z}_p)$, while $I^{\bar{m}} H_3(Y; \mathbb{Z}_p)$ is generated by the restrictions of the Thom space over α and β , say $\hat{\alpha}, \hat{\beta}$ (each of which is homeomorphic as a space to the one point compactification of $S^1 \times \mathbb{R}^2$ since the bundle is trivial over the complement of the divisor).

Now let $X = S^1 \times S^2 \times SY$, where SY is the suspension of Y . The only singular stratum is $S^1 \times S^2 \times \{N, S\}$, where $\{N, S\}$ represents the north and south poles of the suspension. This is a codimension 5 stratum of an 8-dimensional pseudomanifold, and the link of the stratum is Y . Since Y has vanishing middle dimensional middle perversity intersection homology over \mathbb{Z}_p , X is a \mathbb{Z}_p -Witt space, but the middle intersection homology fails to vanish over \mathbb{Q} so that X is not a \mathbb{Q} -Witt space.

Using the formula for the intersection homology of a suspension (see Section 2), together with the Künneth theorem, which holds for intersection homology when one term is a manifold (see [17]), we see that

$$I^{\bar{m}} H_4(X; \mathbb{Z}_p) \cong \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p.$$

If $*$ denotes a basepoint in $S^1 \times S^2$, the generators are $* \times S\hat{\alpha}$, $* \times S\hat{\beta}$, $S^1 \times S^2 \times \alpha$ and $S^1 \times S^2 \times \beta$. If the intersection number $\alpha \cdot \beta = 1$, then the intersection matrix with respect to this basis is

$$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

assuming that SX is thought of as $I \times X / \sim$ (as opposed to $Z \times I / \sim$).

On the other hand,

$$I^{\bar{m}}H_4(X; \mathbb{Q}) \cong \mathbb{Q} \oplus \mathbb{Q} \oplus \mathbb{Q} \oplus \mathbb{Q} \oplus \mathbb{Q},$$

where the first four summands are generated as before and the additional summand is generated by $z = * \times S^2$ times the generator of $I^{\bar{m}}H_2(X; \mathbb{Q})$, which can be represented by T^2 . Its intersection with each generator of $I^{\bar{m}}H_4(X; \mathbb{Q})$, including itself, is 0. For the intersections with $* \times S\hat{\alpha}$, $* \times S\hat{\beta}$, and itself, this can be seen by pushing it off the basepoint in the S^1 direction. For the intersections with $S^1 \times S^2 \times \alpha$ and $S^1 \times S^2 \times \beta$, we can push z off in the direction of the suspension. Thus we obtain a degenerate intersection pairing. The dual of z lives, of course, in $I^{\bar{n}}H_4(X; \mathbb{Q}) \cong \mathbb{Q}^5$, which is generated by our earlier four generators and $S^1 \times * \times ST^2$. With the above convention for suspensions, we have an intersection number $(* \times S^2 \times T^2) \cdot (S^1 \times * \times ST^2) = p$. For $\mathbb{Z}_{p'}$, $p \neq p'$, the computations are the same, replacing all \mathbb{Q} s by $\mathbb{Z}_{p'}$ s.

Similar examples may be obtained easily in higher dimensions. For example, in dimensions $4k$, $k > 2$, we can take the product of X from the previous example with $k - 2$ copies of $\mathbb{C}P^2$. The new space will be Witt (or non-Witt) for exactly the same fields as for X , the middle-dimensional pairing over \mathbb{Z}_p remains nontrivial (and nonsingular), and, if V_i represents the sphere in the i th copy of $\mathbb{C}P^2$ generating the homology in degree 2, then $* \times S^2 \times T^2 \times \prod_{i=1}^{k-2} V_i$ represents a nontrivial \bar{m} -allowable class over \mathbb{Q} and $\mathbb{Z}_{p'}$ whose dual $S^1 \times * \times ST^2 \times \prod_{i=1}^{k-2} V_i$ is \bar{n} -allowable but not \bar{m} -allowable.

Furthermore, applying Remark 3.1 from above, if we carry through the above procedure for a bundle with euler number m , a composite instead of a prime, then we obtain spaces that are \mathbb{Z}_p -Witt for all primes p such that $p \mid m$ but not \mathbb{Q} -Witt nor \mathbb{Z}_p -Witt when $p \nmid m$.

\mathbb{Q} -Witt, but not \mathbb{Z}_p -Witt for some p . We now look for spaces that are \mathbb{Q} -Witt but that fail to be \mathbb{Z}_p -Witt for a single prime or a collection of primes. This corresponds to item (3) of Theorem 1.1. In general, obtaining such spaces is easy; for example, take any even-dimensional closed manifold whose middle homology is all torsion and suspend as many times as desired. However, we would once again like to verify that these actually exhibit the correct existence or lack of middle-dimensional pairings (at least for spaces of dimension $4k$, $k > 1$). It turns out that we can do even this without having to resort to constructions quite as specialized as those in the last section; in particular, we can start with manifolds and introduce a singularity with just a single suspension.

To start off, fix a prime p , and let L be a 3-dimensional lens space with $H_1(L) \cong \mathbb{Z}_p$ (see, e.g. [24, Section 40]). Then we have $H_0(L; \mathbb{Q}) \cong H_3(L; \mathbb{Q}) \cong \mathbb{Q}$, $H_1(L; \mathbb{Q}) = H_2(L; \mathbb{Q}) = 0$, and the same formulas replacing \mathbb{Q} everywhere by $\mathbb{Z}_{p'}$ for $p' \neq p$. However, for coefficients

in \mathbb{Z}_p , $H_i(L; \mathbb{Z}_p) \cong \mathbb{Z}_p$ for $0 \leq i \leq 3$. This follows from the ordinary integer homology of the lens space and the universal coefficient theorem.

If we now let $J = L \times S^1$, then J is a compact orientable 4-manifold with $H_2(J; \mathbb{Z}_p) \cong \mathbb{Z}_p \oplus \mathbb{Z}_p$ and $H_2(J; \mathbb{Q}) = H_2(J; \mathbb{Z}_{p'}) = 0$ for $p' \neq p$. Thus the suspension SJ is \mathbb{Q} -Witt and $\mathbb{Z}_{p'}$ -Witt for all $p' \neq p$, but it is not \mathbb{Z}_p -Witt. By taking products with manifolds, we obtain spaces of all dimensions ≥ 5 with these properties.

Computing with the ordinary Künneth Theorem and the suspension formula for intersection homology (see Section 2, above), we see that

$$I^{\bar{m}}H_i(SJ; \mathbb{Q}) \cong \begin{cases} \mathbb{Q}, & i = 5, \\ \mathbb{Q}, & i = 4, \\ 0, & i = 3, \\ 0, & i = 2, \\ \mathbb{Q}, & i = 1, \\ \mathbb{Q}, & i = 0, \end{cases}$$

and similarly with all \mathbb{Q} s replaced by $\mathbb{Z}_{p'}$. On the other hand,

$$I^{\bar{m}}H_i(SJ; \mathbb{Z}_p) \cong \begin{cases} \mathbb{Z}_p, & i = 5, \\ \mathbb{Z}_p \oplus \mathbb{Z}_p, & i = 4, \\ 0, & i = 3, \\ \mathbb{Z}_p \oplus \mathbb{Z}_p, & i = 2, \\ \mathbb{Z}_p \oplus \mathbb{Z}_p, & i = 1, \\ \mathbb{Z}_p, & i = 0. \end{cases}$$

Now, consider $X = SJ \times S^1 \times S^2$, which is an 8-dimensional pseudomanifold. We have $I^{\bar{m}}H_4(X; \mathbb{Q}) \cong \mathbb{Q} \oplus \mathbb{Q}$, generated by the suspension $S(L \times *_{S^1}) \times *_{S^1 \times S^2}$ and by $(*_L \times S^1) \times S^1 \times S^2$. These cycles are readily checked to be dual to each other. The same is true replacing \mathbb{Q} with $\mathbb{Z}_{p'}$.

On the other hand, let d_i be the i -cell in the standard decomposition of the lens space with one cell in each dimension (see [24]). Then $I^{\bar{m}}H_4(X; \mathbb{Z}_p) \cong \mathbb{Z}_p^6$. The generators are:

$$\begin{array}{ll} S(L \times *_{S^1}) \times *_{S^1 \times S^2} & S(d_2 \times S^1) \times *_{S^1 \times S^2} \\ (d_2 \times *_{S^1}) \times *_{S^1} \times S^2 & (d_1 \times S^1) \times *_{S^1} \times S^2 \\ (d_1 \times *_{S^1}) \times S^1 \times S^2 & (d_0 \times S^1) \times S^1 \times S^2. \end{array}$$

It is the middle row of generators that do not have appropriate duals in $I^{\bar{m}}H_4(X; \mathbb{Z}_p)$. Their duals should be, respectively, $S(d_1 \times S^1) \times S^1 \times *_{S^2}$ and $S(d_2 \times *_{S^1}) \times S^1 \times *_{S^2}$, which, of

course, are generators of $I^{\bar{n}}H_4(X; \mathbb{Z}_p)$. One readily checks geometrically that the intersection numbers of $(d_2 \times *_{S^1}) \times *_{S^1} \times S^2$ and $(d_1 \times S^1) \times *_{S^1} \times S^2$ are 0 with all other generators of $I^{\bar{n}}H_4(X; \mathbb{Z}_p)$ - with the first two and the middle two by pushing off in the $* \times S^1 \times *$ direction and with the last two by pushing up or down in the direction of the suspension.

From here, we may once again obtain examples in all dimensions $4k$, $k > 2$, by taking products with $\mathbb{C}P^2$ s. Also, by taking connected sums with spaces constructed in the exact same way but for different primes in a set $P = \{p_i\}$, we obtain spaces that are \mathbb{Q} -Witt and \mathbb{Z}_p -Witt for any $p \notin P$ but that are not \mathbb{Z}_p -Witt for any $p \in P$.

Remaining questions. We leave the following as open questions:

Question. Are there spaces that are \mathbb{Q} -Witt but that are not \mathbb{Z}_p -Witt for an *infinite* set of primes? Are there spaces that are \mathbb{Q} -Witt but that are not \mathbb{Z}_p -Witt for all but a finite set of primes?

There can be no such compact example with all links compact *manifolds*, since the middle dimensional homology groups would be finitely generated and thus not capable of carrying infinite different types of torsion.

Question. Are there spaces that are not \mathbb{Q} -Witt but that are \mathbb{Z}_p -Witt for an *infinite* set of primes? Are there spaces that are not \mathbb{Q} -Witt but that are \mathbb{Z}_p -Witt for all but a finite set of primes?

4 K -Witt bordism groups

In this section, we discuss the adaptation of Siegel's theorem on \mathbb{Q} -Witt bordism to other coefficient fields. In [27], Siegel notes that, as a consequence of the Poincaré duality on \mathbb{Q} -Witt spaces, for each $k > 0$ there is a well-defined homomorphism from the Witt bordism group $\Omega_{4k}^{\mathbb{Q}\text{-Witt}}$ of compact $4k$ -dimensional \mathbb{Q} -Witt spaces to the Witt group $W(\mathbb{Q})$ of nondegenerate symmetric \mathbb{Q} -bilinear forms, given by taking a \mathbb{Q} -Witt space to its middle dimensional middle-perversity intersection form. One of the principal results of [27] is that this homomorphism is, in fact, an isomorphism and that these bordism groups are 0 in all other dimensions except for $k = 0$, which has $\Omega_0^{\mathbb{Q}\text{-Witt}} \cong \mathbb{Z}$. It then follows from work of Sullivan that, as a homology theory, \mathbb{Q} -Witt bordism, $\Omega^{\mathbb{Q}\text{-Witt}}(\cdot)$, is equivalent to $KO[1/2](\cdot)$. In this section, we extend Siegel's results by computing the K -Witt bordism groups for an arbitrary field K . We prove Theorems 1.2 and 1.3, stated in the introduction. The reader can find more background on Witt groups in [22, 19].

In Subsection 4.1, we provide the basic definitions and some preliminary observations. In Subsection 4.2, we show that the Witt bordism groups (in fact the property of being a K -Witt space) depends only on the characteristic of K . In Subsection 4.3, we prove the analogue of Siegel's theorem, $\Omega_{4k}^{\mathbb{Z}_p\text{-Witt}} \cong W(\mathbb{Z}_p)$ for $k > 0$. In Subsection 4.4, we examine more closely the map $\Omega_{4k}^{\mathbb{F}_q\text{-Witt}} \rightarrow W(\mathbb{F}_q)$ for the finite fields \mathbb{F}_q . Finally, in Subsection 4.5, we show that, as a homology theory, \mathbb{Z}_p -Witt bordism splits into a sum of (shifted) ordinary homology groups with coefficients.

4.1 Preliminaries

Let K be a field. A PL space X is a K -Witt space with boundary if ∂X and $X - \partial X$ are PL pseudomanifolds that satisfy the K -Witt condition and ∂X has a collar in X . This collared boundary requirement is a more restrictive condition than allowing codimension 1 strata in general (which are referred to as “pseudoboundaries” in [10]). We let $\Omega_n^{K\text{-Witt}}$ denote the group of bordism classes of n -dimensional K -Witt spaces, in which X is trivial if X is the boundary of an $n + 1$ dimensional K -Witt space with boundary. See [27, 12] for more details in the \mathbb{Q} -Witt case.

The main invariant of the Witt bordism groups comes from the middle-dimensional intersection pairings. For a $2k$ -dimensional compact oriented K -Witt space, there is a nondegenerate $(-1)^k$ -symmetric intersection pairing $I^{\bar{m}}H_k(X; K) \otimes I^{\bar{m}}H_k(X; K) \rightarrow K$; see [13, 8] for more on the intersection pairing. For $n \equiv 0 \pmod{4}$, the resulting homomorphism $w = w_K : \Omega_n^{K\text{-Witt}} \rightarrow W(K)$ is well-defined. The proof of this in the K case is exactly the same as that for \mathbb{Q} -Witt spaces given in [27, Theorem 2.1], which itself uses intersection homology Poincaré duality to show that the intersection form of a boundary must have a self-annihilating subspace of half the dimension of $I^{\bar{m}}H_k(X; K)$. The basic idea is exactly the same as the proof of signature invariance under manifold bordism. Any pairing with a self-annihilating subspace of half the dimension is trivial in the Witt group; see [22].

4.2 Reduction to prime fields

First, we reduce the problem of computing $\Omega_n^{K\text{-Witt}}$ to the special cases where $K = \mathbb{Z}_p$ or \mathbb{Q} by showing that whether or not X is a K -Witt space is determined entirely by the characteristic of K . For ease of treating all cases simultaneously, we define $\mathbb{Z}_0 := \mathbb{Q}$. We state the key results and then provide the proofs at the end of the subsection.

Lemma 4.1. *Let X be a PL stratified pseudomanifold. Suppose K is a field of characteristic p (which may be 0). Then, (letting $\mathbb{Z}_0 = \mathbb{Q}$), there is a chain isomorphism*

$$I^{\bar{p}}C_*(X; \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} K \rightarrow I^{\bar{p}}C_*(X; K).$$

Corollary 4.2. *Let X be a PL stratified pseudomanifold and K a field of characteristic p , possibly with $p = 0$. Then X is K -Witt if and only if X is \mathbb{Z}_p -Witt (taking $\mathbb{Z}_0 = \mathbb{Q}$).*

Corollary 4.3. *If K is of characteristic p , $\Omega_n^{K\text{-Witt}} = \Omega_n^{\mathbb{Z}_p\text{-Witt}}$. In particular, if K has characteristic 0, $\Omega_n^{K\text{-Witt}} = \Omega_n^{\mathbb{Q}\text{-Witt}}$.*

Corollary 4.3 follows immediately from Corollary 4.2.

To see how this reduction relates to the Witt group invariants, note that for any inclusion of fields $\mathbb{Z}_p \hookrightarrow K$ (or $\mathbb{Q} \hookrightarrow K$ if $p = 0$), we obtain a commutative diagram

$$\begin{array}{ccc} \Omega_{4k}^{\mathbb{Z}_p\text{-Witt}} & \xrightarrow{w_{\mathbb{Z}_p}} & W(\mathbb{Z}_p) \\ \downarrow = & & \downarrow \\ \Omega_{4k}^{K\text{-Witt}} & \xrightarrow{w_K} & W(K). \end{array} \tag{1}$$

Here the righthand vertical map is induced by the field homomorphism $\mathbb{Z}_p \hookrightarrow K$. To see that this diagram commutes, first note that by Lemma 4.1, $I^{\bar{m}}H_{2k}(X; K)$ is generated over K by elements of the form $[\xi] \otimes 1$, where $\xi \in I^{\bar{m}}H_{2k}(X; \mathbb{Z}_p)$. So, if X^{4k} is K -Witt (and hence \mathbb{Z}_p -Witt) and we choose a basis for $I^{\bar{m}}H_{2k}(X; \mathbb{Z}_p)$, then the resulting intersection pairing matrix for the dual K pairing on $I^{\bar{m}}H_{2k}(X; K)$ is identical to the intersection pairing matrix for the \mathbb{Z}_p pairing on $I^{\bar{m}}H_{2k}(X; \mathbb{Z}_p)$. Thus the same matrix with entries in \mathbb{Z}_p represents both $w_{\mathbb{Z}_p}(X) \in W(\mathbb{Z}_p)$ and $w_K(X) \in W(K)$. This is consistent with the map $W(\mathbb{Z}_p) \rightarrow W(K)$ induced by inclusion.

It follows that w_K is determined entirely by $w_{\mathbb{Z}_p}$, which will be studied in the next subsection for $p \neq 0$.

We return now to the deferred proofs.

Proof of Lemma 4.1. Define $\Phi : I^{\bar{p}}C_*(X; \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} K \rightarrow I^{\bar{p}}C_*(X; K)$ by $\Phi(\sum_i \xi_i \otimes k_i) = \sum_i k_i \xi_i$, where each $\xi_i \in I^{\bar{p}}C_*(X; \mathbb{Z}_p)$ and $k_i \in K$. Note that $k_i \xi_i$ makes sense as each ξ_i equals $\sum m_j \sigma_j$ for $m_j \in \mathbb{Z}_p$ and σ_j a simplex of some triangulation of X , and $k_i m_j$ makes sense as an element of K . It is easy to check that Φ is a chain map and well-defined.

Perhaps the simplest way to check that Φ is injective is to consider the following commutative diagram:

$$\begin{array}{ccc} I^{\bar{p}}C_*(X; \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} K & \xrightarrow{\Phi} & I^{\bar{p}}C_*(X; K) \\ \downarrow & & \downarrow \\ C_*(X; \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} K & \xlongequal{\quad} & C_*(X; K). \end{array}$$

Note that it is clear from the definitions that $I^{\bar{p}}C_*(X; G) \rightarrow C_*(X; G)$ is always an inclusion. Since the tensor product over a field is left exact, the vertical arrows are inclusions, and the bottom map is a standard isomorphism. It follows from the diagram that Φ is injective.

For surjectivity, let $\xi = \sum k_i \sigma_i \in I^{\bar{p}}C_*(X; K)$. Note that the sum is finite, since each element of $C_*(X)$ lives in a fixed triangulation. Consider the \mathbb{Z}_p vector subspace V of K spanned by the k_i . Let $\{x_j\}_{j=1}^N$ be a basis for V . Then each $k_i = \sum n_j x_j$, $n_j \in \mathbb{Z}_p$. Using this, we can rewrite ξ in the form $\sum x_j \xi_j$, where $\xi_j \in C_*(X; \mathbb{Z}_p)$. We claim that each $\xi_j \in I^{\bar{p}}C_*(X; \mathbb{Z}_p)$, from which it will follow that $\xi = \Phi(\sum \xi_j \otimes x_j)$. It is clear that each simplex σ appearing in each ξ_j must be allowable since each occurs with non-zero coefficient in ξ . The point is to show that each $\partial \xi_j$ is allowable, which is not immediately clear. However, suppose that τ is a simplex that appears with nonzero (\mathbb{Z}_p -)coefficient in $\partial \xi_j$ for some j . The total coefficient of τ in $\partial \xi$ must have the form $\sum_j x_j m_j$, where m_j is the coefficient of τ in $\partial \xi_j$. Since τ appears nontrivially in $\partial \xi_j$ for some j , some $m_j \not\equiv 0 \pmod{p}$, and so $\sum_j x_j m_j \neq 0$, as the ξ_j are linearly independent as a vector space basis. Thus τ appears nontrivially in $\partial \xi$, and hence must be allowable, because ξ is an allowable chain. \square

Remark 4.4. This lemma can be shown more generally over topological pseudomanifolds using the sheaf approach to intersection homology. We provide a PL chain level proof, more in keeping with the spirit of the current paper.

Proof of Corollary 4.2. From the lemma and the algebraic universal coefficient theorem for fields, for a compact PL pseudomanifold, $I^{\bar{p}}H_*(X; K) \cong I^{\bar{p}}H_*(X; \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} K$. So, for a compact link L , $I^{\bar{p}}H_i(L; K)$ vanishes if and only if $I^{\bar{p}}H_i(L; \mathbb{Z}_p)$ vanishes. The corollary follows. \square

4.3 Witt bordism over \mathbb{Z}_p

In this section we compute $\Omega_n^{\mathbb{Z}_p\text{-Witt}}$. For a further discussion of the case of more general finite fields, see Section 4.4 below.

Theorem 4.5. *Let $p \neq 0$ be a prime.*

1. $\Omega_0^{\mathbb{Z}_p\text{-Witt}} \cong \mathbb{Z}$.
2. For $n \not\equiv 0 \pmod{4}$, $\Omega_n^{\mathbb{Z}_p\text{-Witt}} = 0$.
3. For $n \equiv 0 \pmod{4}$, $n > 0$, the homomorphism $w : \Omega_n^{\mathbb{Z}_p\text{-Witt}} \rightarrow W(\mathbb{Z}_p)$ that assigns to X^n the intersection form on $I^{\bar{m}}H_{n/2}(X; \mathbb{Z}_p)$ is an isomorphism.

We can observe immediately that all 0- and 1-dimensional \mathbb{Z}_p -Witt spaces are, of course, manifolds, so that $\Omega_0^{\mathbb{Z}_p\text{-Witt}} \cong \mathbb{Z}$, as for manifolds and \mathbb{Q} -Witt spaces. Furthermore, as for \mathbb{Q} -Witt spaces, $\Omega_{2k+1}^{\mathbb{Z}_p\text{-Witt}} = 0$, since if X is an odd-dimensional \mathbb{Z}_p -Witt space, then the closed cone $\bar{c}X$ is also \mathbb{Z}_p -Witt, as the new stratum consisting of the cone vertex has even codimension.

This leaves the Witt spaces of positive even dimension.

We must next show the following:

Proposition 4.6. *$w : \Omega_n^{\mathbb{Z}_p\text{-Witt}} \rightarrow W(\mathbb{Z}_p)$ is an isomorphism for all primes p and $n = 4k > 0$, and $\Omega_n^{\mathbb{Z}_p\text{-Witt}} = 0$ for $n \equiv 2 \pmod{4}$.*

Proof of Proposition 4.6. The proof is mostly analogous to that of Siegel's theorem for \mathbb{Q} -Witt bordism. The main difference is the proof of surjectivity for $0 \neq q \equiv 0 \pmod{4}$. We will demonstrate this surjectivity and then discuss the one significant change from Siegel's proof of injectivity that we must make in the \mathbb{Z}_p situation. The proof that $\Omega_q^{\mathbb{Z}_p\text{-Witt}} = 0$ for $q \equiv 2 \pmod{4}$ is included with the proof of injectivity of w for $0 < q \equiv 0 \pmod{4}$.

The surjectivity of $w : \Omega_{4k}^{\mathbb{Z}_p\text{-Witt}} \rightarrow W(\mathbb{Z}_p)$ is contained in the following proposition.

Proposition 4.7. *Given any element of $x \in W(\mathbb{Z}_p)$ and any integer $k > 0$, there exists a $4k$ -dimensional \mathbb{Z}_p -Witt space X whose intersection pairing on $I^{\bar{m}}H_{2k}(X; \mathbb{Z}_p)$ represents x .*

Proof. It follows from the theory of Witt rings (see [19, Chapter 2]) that for a finite field F , the Witt ring $W(F)$ is isomorphic to a semi-direct product $Q(F) = \mathbb{Z}_2 \ltimes \dot{F}/\dot{F}^2$, where \dot{F} is the multiplicative group $F - \{0\}$ and $\dot{F}/\dot{F}^2 \cong \mathbb{Z}_2$. The group operation in $Q(F)$ is given by $(e, f) \cdot (e', f') = (e + e', (-1)^{ee'} ff')$, and the isomorphism $W(F) \rightarrow Q(F)$ is given by the pair of operators (\dim_0, d_{\pm}) . Here \dim_0 takes a representative symmetric bilinear form to its dimension $\pmod{2}$ and d_{\pm} takes a representative form of dimension n to $(-1)^{n(n-1)/2}$ times the determinant of the matrix representing the form.

When $F = \mathbb{F}_q$ with $q \equiv 3 \pmod{4}$, then $W(F) \cong Q(F) \cong \mathbb{Z}_4$. The generator is $(1, 1)$ and $Q(F) = \{(0, 1), (1, 1), (0, -1), (1, 0)\}$ (note that -1 is not a square when $q \equiv 3 \pmod{4}$). In terms of forms, $W(F)$ is therefore generated by $\langle 1 \rangle$.

If $q \equiv 1 \pmod{4}$, then $Q(F)$ splits as $\mathbb{Z}_2 \times \mathbb{Z}_2$. If $s \in \mathbb{F}_q$ is not a square, then the elements of $Q(F)$ are represented by $\{(0, 1), (1, 1), (1, s), (0, s)\}$, generated in $W(F)$ by $\langle 1 \rangle$ and $\langle s \rangle$.

If q is even, then $W(\mathbb{F}_q) \cong \mathbb{Z}_2$ [22].

So when $p \equiv 3 \pmod{4}$, it is easy to find manifolds of dimension $4k$, $k \geq 1$ whose middle dimensional pairings represent any element of the Witt group. Specifically, the pairing on $\mathbb{C}P^{2k}$ is the generator of $W(\mathbb{Z}_p)$ and taking connected sums of $\mathbb{C}P^k$ s yields representatives of any element of $W(\mathbb{Z}_p)$.

For $p \equiv 1 \pmod{4}$, $\mathbb{C}P^{2k}$ again yields the generator $\langle 1 \rangle$ of the Witt group $W(\mathbb{Z}_p)$. To obtain the generator $\langle s \rangle$, we can proceed with Thom spaces as in Section 3.1. If $s > 0$ represents a non-square unit in \mathbb{Z}_p , let U be the normal bundle of the connected sum of s copies of the generator $[V_i] \in H_{2k}(\mathbb{C}P^{2k})$, represented by embedded spheres S_i^{2k} , in the connected sum of s copies of $\mathbb{C}P^{2k}$.⁴ Let \hat{U} be the associated Thom space. Then, as in the computations in Section 3.1, $I^{\bar{m}}H_{2k}(\hat{U}; \mathbb{Z}_p) \cong \text{Im}: (H_{2k}(S^{2n}; \mathbb{Z}_p) \cong \mathbb{Z}_p \xrightarrow{\cap \epsilon} H_0(S^{2k}; \mathbb{Z}_p) \cong \mathbb{Z}_p)$. Since the euler number of the bundle is the unit s , $I^{\bar{m}}H_{2k}(\hat{U}; \mathbb{Z}_p) \cong \mathbb{Z}_p$ generated by $[V] = \sum [V_i] = [S^{2n}]$. Furthermore, $[V] \cdot [V] = s$. Thus, since the only singularity of \hat{U} has codimension $4k$, \hat{U} is a Witt space with intersection form $\langle s \rangle$. So we have constructed spaces representing both generators of $W(\mathbb{Z}_p)$.

For $p = 2$, $W(\mathbb{Z}_p)$ is generated simply by $\langle 1 \rangle$, and we can use again any $\mathbb{C}P^{2k}$ as a geometric realization. \square

Turning to the injectivity of $w : \Omega_{4k}^{\mathbb{Z}_p\text{-Witt}} \rightarrow W(\mathbb{Z}_p)$, as well as the fact that $\Omega_n^{\mathbb{Z}_p\text{-Witt}} = 0$ for $n \equiv 2 \pmod{4}$, we note again that the proof is nearly identical to that of the \mathbb{Q} -Witt case, though some care must be taken, primarily with the use of geometric cycles (see the discussion at the end of Section 4.4 for more elaboration on what can go wrong over fields more general than \mathbb{Z}_p). We discuss this issue and refer the reader to [27] for the remainder of the proof.

Siegel's proof of injectivity over \mathbb{Q} begins by supposing we have a \mathbb{Q} -Witt space X^{2k} with an isotropic element $[z] \in I^{\bar{m}}H_k(X; \mathbb{Q})$, i.e. $[z] \cdot [z] = 0$. Siegel then finds a representative of $[z]$ by an *irreducible cycle* z , meaning that $H_k(|z|; \mathbb{Z}) \cong \mathbb{Z}$ and such that the generator of this homology group has coefficient ± 1 on every k simplex of $|z|$ in some triangulation of $|z|$. The key point is that the support $|z|$ of z should have infinite cyclic k th homology, and it should be generated by $|z|$, itself, considered as the cycle represented by its fundamental class.

In the \mathbb{Z}_p case, the construction of z should be altered slightly. Given any $2k$ -dimensional Witt space, any class $[z] \in I^{\bar{m}}H_k(X; \mathbb{Z}_p)$ will satisfy $[z] \cdot [z] = 0$ if $2k \equiv 2 \pmod{4}$, while if $2k \equiv 0 \pmod{4}$ and $w(X) = 0 \in W(\mathbb{Z}_p)$, such a cycle certainly exists. We construct a " \mathbb{Z}_p -irreducible" representative cycle z for $[z]$ by slightly modifying Siegel's construction to obtain a z such that $H_k(|z|; \mathbb{Z}_p) \cong \mathbb{Z}_p$, generated by a "fundamental class" of $|z|$. The

⁴Note that here it is crucial that we work in \mathbb{Z}_p and not some \mathbb{F}_{p^n} so that s can be represented by an integer.

quotation marks indicate that this is not quite the right language since $|z|$ might have a codimension one singularity - however since z is a cycle, these singularities will cancel when thinking of z as a chain, and so the idea of a fundamental class makes some sense.

Briefly, we make z irreducible as follows: Choose an arbitrary representative y for $[z]$ in some triangulation of X . Then $y = \sum n_i \sigma_i$, where $n_i \in \mathbb{Z}_p$ and each σ_i is a unique oriented k -simplex. Choose $m_i \equiv n_i \pmod p$ such that $0 \leq m_i < p$, and abusing notation, let $y = \sum m_i \sigma_i$. Note that the interior of each k -simplex of y lies in $X - \Sigma$ due to the allowability conditions for an intersection chain. Now, for each σ_i such that $m_i \neq 0$, we may use a relative stratified general position argument following McCrory [21] (see also [8]) to PL isotope the interiors of the m_i copies of σ_i , rel boundary, into stratified general position with respect to each other, and in such a way that none of the new m_i copies of σ_i intersect any of the m_j copies of σ_j similarly created, except along boundaries. This is more or less an alternative description of Step 1 of the proof of Siegel's [27, Lemma 2.2], except that Siegel separates by isotopy entire open j -strata of $|y|$. Arguments along the lines of the stratified homotopy invariance of intersection homology (see [9]) show that this new chain, \bar{y} , also represents $[z]$, and clearly $\partial \bar{y} = \partial y = 0 \in C_{k-1}(X; \mathbb{Z}_p)$. Furthermore, y has the form $y = \sum \sigma_i$, where the sum is taken over those oriented simplices in the support of y . To form z , one then connects all of the k -simplices of y by orientation respecting pipes; see [27, page 1087] for more details. Then one also has z of the form $z = \sum \sigma_i$ (with different σ s from \bar{y}), and clearly z generates $H_k(|z|; \mathbb{Z}_p) \cong \mathbb{Z}_p$.

This z can then be used in the remainder of a \mathbb{Z}_p analogue of Siegel's injectivity proof. □

This completes the proof of Theorem 4.5. □

Remark 4.8. Incidentally, the process of the injectivity proof can be used to find many more examples of \mathbb{Z}_p -Witt spaces with vanishing middle-dimensional $I^m H_k$ since, by copying the proof of Siegel's [27, Proposition III.3.1] exactly, $I^m H_k(\hat{U}; \mathbb{Z}_p) = 0$, where U is a regular neighborhood of our irreducible z (still assuming $[z] \cdot [z] = 0$), and $\hat{U} = U \cup \bar{c}(\partial U)$.

4.4 Witt bordism over finite fields

In this section, we observe the ramifications of the results of the previous section to $\Omega_n^{\mathbb{F}_q\text{-Witt}}$ for the finite field \mathbb{F}_q . We also provide an illustrative example.

Proposition 4.9. *For $n \equiv 0 \pmod 4$, $n > 0$, the homomorphism $w : \Omega_n^{\mathbb{F}_q\text{-Witt}} \rightarrow W(\mathbb{F}_q)$ that assigns to X^n the intersection form on $I^m H_{n/2}(X; \mathbb{F}_p)$ is an isomorphism, except when $q = p^m$, $p \equiv 3 \pmod 4$, m even. In these exceptional cases, $\Omega_n^{\mathbb{F}_p\text{-Witt}}$ is isomorphic to $W(\mathbb{Z}_p)$, and w is the homomorphism induced by inclusion $W(\mathbb{Z}_p) \rightarrow W(\mathbb{F}_p)$.*

The proposition follows from Theorem 4.5, Corollary 4.3, and the commutativity of diagram (1) for $k, m > 0$. In particular, Corollary 4.3 tells us that the map $w : \Omega_{4k}^{\mathbb{Z}_p\text{-Witt}} \rightarrow W(\mathbb{Z}_p)$ is an isomorphism, and so from the commutativity of the diagram, $w : \Omega_{4k}^{\mathbb{F}_p\text{-Witt}} \rightarrow W(\mathbb{F}_p)$ is isomorphic to the homomorphism $W(\mathbb{Z}_p) \rightarrow W(\mathbb{F}_p)$. We recall the properties of this homomorphism.

Consider the homomorphism $\psi : W(\mathbb{Z}_p) \hookrightarrow W(\mathbb{F}_{p^m})$ induced by the natural inclusion $\mathbb{Z}_p \hookrightarrow \mathbb{F}_{p^m}$. If $p \equiv 1 \pmod{4}$, then $p^m \equiv 1 \pmod{4}$ for all $m > 0$, while if $p \equiv 3 \pmod{4}$, $p^{2m+1} \equiv 3 \pmod{4}$ and $p^{2m} \equiv 1 \pmod{4}$. In all cases except the last (p^{2m} for $p \equiv 3 \pmod{4}$), ψ is an isomorphism. This follows from the discussion in the proof of Proposition 4.7: in these cases we know that $W(\mathbb{Z}_p) \cong W(\mathbb{F}_{p^m})$ abstractly and ψ clearly preserves dimension and determinant of the form, which constitute a complete set of invariants.

In the exceptional case, $W(\mathbb{Z}_p) \cong \mathbb{Z}_4$, generated by $\langle 1 \rangle$, but $W(\mathbb{F}_{p^{2m}}) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$, in which $\langle 1 \rangle$ is one of the generators. So ψ maps $W(\mathbb{Z}_p)$ onto one of the summands of $W(\mathbb{F}_{p^m})$ with kernel \mathbb{Z}_2 .

What does this mean geometrically? As an example, consider \mathbb{Z}_3 and $\mathbb{F}_9 \cong \mathbb{Z}_3[x]/\langle x^2-2 \rangle$. Let $\{1, x\}$ denote a basis of \mathbb{F}_9 as a 2-dimensional vector space over \mathbb{Z}_3 . We showed above that $\mathbb{C}P^2$ is a generator of $\Omega_4^{\mathbb{Z}_3\text{-Witt}}$ corresponding to $\langle 1 \rangle$ in $W(\mathbb{Z}_3)$. According to the results of the preceding section, $\Omega_4^{\mathbb{Z}_3\text{-Witt}} \cong W(\mathbb{Z}_3) \cong \mathbb{Z}_4$ so that $\mathbb{C}P^2$ has order 4. Since $\Omega_4^{\mathbb{Z}_3\text{-Witt}} = \Omega_4^{\mathbb{F}_9\text{-Witt}}$, $\mathbb{C}P^2$ also has order 4 in the latter group. However $W(\mathbb{F}_9) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$, so, in particular, $w(\mathbb{C}P^2 \# \mathbb{C}P^2) = 0 \in W(\mathbb{F}_9)$. To see where a proof of injectivity along the lines of Siegel breaks down, observe that $w(\mathbb{C}P^2 \# \mathbb{C}P^2)$ is represented by $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. Since this pairing is nonzero in $W(\mathbb{Z}_3)$, in order to find an isotropic element over \mathbb{F}_9 , we must mix 1 and x nontrivially. For example, let V_1 and V_2 denote two disjoint spheres in $\mathbb{C}P^2 \# \mathbb{C}P^2$ generating $I^{\bar{m}} H_2(\mathbb{C}P^2 \# \mathbb{C}P^2; \mathbb{F}_9) \cong H_2(\mathbb{C}P^2 \# \mathbb{C}P^2; \mathbb{F}_9) \cong H_2(\mathbb{C}P^2 \# \mathbb{C}P^2) \otimes \mathbb{F}_9$. An isotropic element is given by $[V_1] + x[V_2]$, since $([V_1] + x[V_2]) \cdot ([V_1] + x[V_2]) = 1 + x^2 = 1 + 2 = 0 \in \mathbb{F}_9$. There is no way to create a cycle representing $[V_1] + x[V_2]$ that is irreducible in the sense discussed in the injectivity proof in Section 4.3, since the mixing of 1 and x coefficients would prevent us from piping simplices together (for that matter, we cannot make sense of taking x copies of something).

4.5 \mathbb{Z}_p -Witt bordism as a generalized homology theory

This subsection contains a computation of the homology theory $\Omega_*^{\mathbb{Z}_p\text{-Witt}}(\cdot)$. As for rational Witt bordism, K -Witt bordism yields a generalized homology theory for any K ; as noted by Siegel [27, Chapter IV], this follows from Akin [1, Proposition 7], making the obvious generalizations from unoriented to oriented bordism. Akin's axioms are easy to check for K -Witt spaces with boundary, by making use of the collars on the boundaries to see that Akin's cuttings and pastings do not create new links that would violate the Witt conditions.

It follows from the results of the preceding sections that the only parameter that matters in K -Witt-bordism is the characteristic of the field K . By Siegel, $\Omega_*^{\mathbb{Q}\text{-Witt}}(\cdot) \cong KO[1/2]_*(\cdot)$, so we focus on fields of finite characteristic. In particular, there is no loss of generality limiting ourselves to \mathbb{Z}_p . It turns out that $\Omega_*^{\mathbb{Z}_p\text{-Witt}}(\cdot)$ splits as a sum of ordinary homology theories with coefficients, which is the content of the following theorem.

Theorem 4.10. *As a homology theory on CW complexes, \mathbb{Z}_p -Witt bordism splits as a direct sum of ordinary homology with coefficients. In particular,*

$$\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X) \cong \bigoplus_{r+s=n} H_r(X; \Omega_s^{\mathbb{Z}_p\text{-Witt}}).$$

Proof. Brown representability tells us that \mathbb{Z}_p -Witt bordism is representable by a spectrum; see, e.g. [28, Theorem 14.35, Corollary 14.36, and Remark 1 on page 331]. To see that \mathbb{Z}_p -Witt bordism on CW complexes satisfies the wedge axiom, and hence the conditions in [28, Remark 1], note the unreduced version of the wedge axiom within the proof of [28, Proposition 10.16] and that our bordism theory breaks into a sum over what happens on connected components. We let W_p denote the spectrum yielding \mathbb{Z}_p -Witt bordism. We will show below that W_p is, in fact, an MSO -module spectrum, but the proof of this fact is deferred for now.

Let X be a compact CW complex so that the homology of X is finitely generated. We consider the Atiyah-Hirzebruch spectral sequence for $\Omega_*^{\mathbb{Z}_p\text{-Witt}}(X)$, whose $E_{r,s}^2$ term is $H_r(X; \Omega_s^{\mathbb{Z}_p\text{-Witt}})$. Since $\Omega_s^{\mathbb{Z}_p\text{-Witt}}$ is 2-primary for all $s > 0$, the only odd torsion in the spectral sequence is in the terms $H_r(X; \Omega_0^{\mathbb{Z}_p\text{-Witt}}) \cong H_r(X; \mathbb{Z})$, which lie along the x -axis. It follows that the odd torsion part of the spectral sequence splits off and that the odd torsion subgroup of $\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X)$ is isomorphic to the odd torsion subgroup of $H_n(X)$.

Now, for the rest of $\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X)$, we consider instead the theory $\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X; \mathbb{Z}_{(2)}) \cong \Omega_n^{\mathbb{Z}_p\text{-Witt}}(X) \otimes \mathbb{Z}_{(2)}$ with 2-local coefficients (see [26, Section II.5] for a general reference). In this case, the Atiyah-Hirzebruch E^2 terms look like $H_r(X; \Omega_s^{\mathbb{Z}_p\text{-Witt}} \otimes \mathbb{Z}_{(2)})$. Since $\Omega_s^{\mathbb{Z}_p\text{-Witt}}$ is 2-torsion for $s > 0$, in this case the terms are identical to those from the spectral sequence before the localization, while the $(r, 0)$ term becomes $H_r(X; \mathbb{Z}_{(2)})$. Now, this spectral sequence degenerates at the E^2 term by a theorem of Taylor and Williams [29, Section 2] that states that any 2-local MSO -module spectrum (in this case $(W_p)_{(2)}$) splits as a wedge of Eilenberg-MacLane spectra. The upshot of this is the degeneration of the Atiyah-Hirzebruch spectral sequence. By this degeneration,

$$\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X) \otimes \mathbb{Z}_{(2)} \cong \Omega_n^{\mathbb{Z}_p\text{-Witt}}(X; \mathbb{Z}_{(2)}) \cong H_n(X; \mathbb{Z}_{(2)}) \oplus \bigoplus_{\substack{r+s=n \\ s>0}} H_r(X; \Omega_s^{\mathbb{Z}_p\text{-Witt}}).$$

The group $\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X)$ must be finitely generated (since everything else in the spectral sequence is), so, by basic facts about localization, we can read off the 2-primary and infinite cyclic summands of $\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X)$ from $\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X) \otimes \mathbb{Z}_{(2)}$. In particular, any 2-primary components are shared between the two, and \mathbb{Z} summands of the former, correspond bijectively with $\mathbb{Z}_{(2)}$ terms in the latter, which can come only from the summand $H_n(X; \mathbb{Z}_{(2)})$. Thus, modulo odd torsion, we must have $\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X) \cong H_n(X; \mathbb{Z}) \oplus \bigoplus_{\substack{r+s=n \\ s>0}} H_r(X; \Omega_s^{\mathbb{Z}_p\text{-Witt}})$.

But now we know that the odd torsion of $\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X)$ is that of $H_n(X)$. So, putting everything together, we must have

$$\Omega_n^{\mathbb{Z}_p\text{-Witt}}(X) \cong H_n(X; \mathbb{Z}) \oplus \bigoplus_{\substack{r+s=n \\ s>0}} H_r(X; \Omega_s^{\mathbb{Z}_p\text{-Witt}}) \cong \bigoplus_{r+s=n} H_r(X; \Omega_s^{\mathbb{Z}_p\text{-Witt}}),$$

since $\Omega_0^{\mathbb{Z}_p\text{-Witt}} \cong \mathbb{Z}$.

For infinite CW complexes, we can now use the preceding formula and take direct limits (see [28, page 331, Remark 1]). \square

Now we return to the following lemma.

Lemma 4.11. W_p is an MSO -module spectrum.

Proof. To begin with, W_p is, in the language of Rudyak [26, Definition III.7.7], a quasi-module spectrum over the Thom spectrum MSO . This means that for CW pairs (X, A) and (Y, B) and letting $\Omega_*^{SO}(\cdot)$ be smooth oriented bordism, there is a pairing $\Omega_*^{SO}(X, A) \otimes \Omega_*^{\mathbb{Z}_p\text{-Witt}}(Y, B) \rightarrow \Omega_*^{\mathbb{Z}_p\text{-Witt}}(X \times Y, X \times B \cup A \times Y)$ that possesses the expected associativity. To see that such a pairing exists, we observe that smooth oriented manifolds are all \mathbb{Z}_p -Witt spaces for any p and so is the product of a smooth oriented manifold with a \mathbb{Z}_p -Witt space (the links don't change). Furthermore, we should check that if M^m is a manifold with boundary and N^n is a \mathbb{Z}_p -Witt space with \mathbb{Z}_p -Witt boundary, then $\partial(M \times N) = M \times \partial N \cup_{\partial M \times \partial N} \partial M \times N$ is \mathbb{Z}_p -Witt. The links in $\text{int}(M) \times \partial N$ are the same as the links of ∂N , and the links of $\partial M \times N$ are the same as the links of N . Now, if $x \in \partial M \times \partial N$, then x has a distinguished neighborhood in this space of the form $\mathbb{R}^{m-1} \times \mathbb{R}^{n-1-k} \times cL$. In $M \times \partial N$, this neighborhood expands to the form $\mathbb{R}^{m-1} \times \mathbb{R}^{n-1-k} \times cL \times (-1, 0]$, where the $(-1, 0]$ coordinate represents the collar of ∂M in M . But in $\partial M \times N$, this neighborhood similarly expands to $\mathbb{R}^{m-1} \times \mathbb{R}^{n-1-k} \times cL \times [0, 1)$. where $[0, 1)$ is the collar of ∂N in N . Putting these together, x has a neighborhood of the form $\mathbb{R}^{m+n-1-k} \times cL$, and so again the link of x in $\partial(M \times N)$ is L , which is a link in ∂N , and hence satisfies the Witt condition.

It is not true in general that being a quasi-module spectrum leads to being an actual module spectrum (see [26]), but we will be able to show this for W_p .

We note that both MSO and W_p have finite \mathbb{Z} -type, meaning that they are bounded below (i.e. their coefficient groups vanish below a certain dimension) and each $\pi_i(MSO)$ and $\pi_i(W_p)$ is finitely generated. For MSO this is well-known; see, e.g., [23, Theorem 18.8]. For W_p , this follows from Theorem 4.5. Thus, by [26, Proposition II.4.26.ii], we may assume, up to equivalence, that MSO and W_p have finite type, i.e. that each skeleton of each spectrum is finite (see [26, Definition II.1.2.e]).

We can now apply a module spectrum version of [26, Theorem III.7.3]. It is noted by Rudyak immediately prior to his [26, Theorem III.7.8] that there is such a module spectrum version of his Theorem III.7.3, which gives conditions for when a quasi-ring spectrum is in fact a ring spectrum. The actual hypotheses of his Theorem III.7.8 as stated are more restrictive than those of Theorem III.7.3 only because this is what is needed in the rest of the book, but a version of Theorem III.7.3 in its full generality can be directly applied to the case of quasi-module spectra. The quasi-module version of this theorem says the following: Since our spectra have finite \mathbb{Z} -type, our MSO -quasi-module structure will be induced by a unique (up to homotopy) MSO -module structure on W_p if the following groups vanish: $\varprojlim^1 \{W_p^{-1}(W_p^{(n)})\}$, $\varprojlim^1 \{W_p^{-1}(MSO^{(n)} \wedge W_p^{(n)})\}$, and $\varprojlim^1 \{W_p^{-1}(MSO^{(n)} \wedge MSO^{(n)} \wedge W_p^{(n)})\}$. To clarify, the superscript -1 stands, in each case, for the degree -1 generalized cohomology group, while $E^{(n)}$ stands for the n -skeleton of the spectrum E . The existence of a module morphism $MSO \wedge W_p \rightarrow W_p$ inducing the module structure on homology is automatic, but the vanishing of these three groups is necessary to yield, respectively, the multiplicative unit, uniqueness, and associativity for the module spectrum structure.

To show that these groups vanish, we consider the Atiyah-Hirzebruch spectral sequences. Let $\mathfrak{X}^{(n)}$ be any of $W_p^{(n)}$, $MSO^{(n)} \wedge W_p^{(n)}$, or $MSO^{(n)} \wedge MSO^{(n)} \wedge W_p^{(n)}$. Then, for $W_p^{-1}(\mathfrak{X}^{(n)})$, the relevant E_2 terms of the spectral sequence will have the form $H^i(\mathfrak{X}^{(n)}; \Omega_{-j}^{\mathbb{Z}_p\text{-Witt}})$, with $i + j = -1$. For $j > 0$, these terms vanish obviously. For $j = 0, i = -1$, they also vanish. To see this, note that since $\Omega_k^{\mathbb{Z}_p\text{-Witt}} = \pi_k(W_p) = 0$ and $\Omega_k^{SO} = \pi_k(MSO) = 0$ for $k < 0$, MSO and W_p are connected spectra (this is just the definition of connected spectrum). By [26, Proposition II.4.5.iv], for any spectrum E , the inclusion $E^{(k)} \subset E$ induces isomorphisms on π_i for $i \leq k$, and by [26, Lemma II.4.2], we may assume (by replacing spectra with equivalent ones) that, if E is connected, $E^{(k)} = *$ for $k \leq -1$. It follows from these two facts that $MSO^{(n)}$ and $W_p^{(n)}$ are also connected spectra. Furthermore, by [26, Proposition II.4.5.i], the smash products of connected spectra are connected. In particular, then, each of our $\mathfrak{X}^{(n)}$ is connected. So, by [26, Corollary II.4.7], $H_i(\mathfrak{X}^{(n)}) = 0$ for $i < 0$, and, by the universal coefficient theorem for spectra with coefficients in an abelian group [26, Theorem II.4.9], $H^i(\mathfrak{X}^{(n)}; G) = 0$ for any abelian group G and any $i \leq -1$.

Finally, we consider the terms $H^i(\mathfrak{X}^{(n)}; \Omega_{-j}^{\mathbb{Z}_p\text{-Witt}})$ for $j < 0, i > -1$. As noted above, we can assume up to equivalence that the skeleta $W_p^{(n)}$ and $MSO^{(n)}$ are finite spectra, and, as a consequence of [26, Proposition II.1.5.ii,iii], the (co)homology of a finite spectrum is the same as the (co)homology of some finite CW complex. Thus, applying the Künneth theorem to the smash products $\mathfrak{X}^{(n)}$ (see the remarks following [26, Theorem II.4.11]), the remaining terms $H^i(\mathfrak{X}^{(n)}; \Omega_{-j}^{\mathbb{Z}_p\text{-Witt}})$, $i > -1$, look like the cohomology of products of finite complexes with coefficients in the finite groups $\Omega_{-j}^{\mathbb{Z}_p\text{-Witt}}$, $j < 0$. Therefore, they must vanish for sufficiently large i and will be finite groups otherwise.

It now follows from the Atiyah-Hirzebruch spectral sequence that each $W_p^{-1}(\mathfrak{X}^{(n)})$ is finite, and this will suffice to make the above \varprojlim^1 groups vanish by [26, Corollary III.2.18], which notes that $\varprojlim^i \mathfrak{A} = 0$ for $i > 0$ if \mathfrak{A} is a system of finite abelian groups. \square

References

- [1] Ethan Akin, *Stiefel-Whitney homology classes and bordism*, Trans. Amer. Math. Soc. **205** (1975), 341–359. MR MR0358829 (50 #11288)
- [2] Markus Banagl, *Extending intersection homology type invariants to non-Witt spaces*, vol. 160, Memoirs of the Amer. Math. Soc., no. 760, American Mathematical Society, Providence, RI, 2002.
- [3] ———, *Topological invariants of stratified spaces*, Springer Monographs in Mathematics, Springer-Verlag, New York, 2006.
- [4] Markus Banagl and Greg Friedman, *Triangulations of 3-dimensional pseudomanifolds with an application to state-sum invariants*, Algebr. Geom. Topol. **4** (2004), 521–542.
- [5] A. Borel et. al., *Intersection cohomology*, Progress in Mathematics, vol. 50, Birkhauser, Boston, 1984.

- [6] Glen Bredon, *Topology and geometry*, Springer-Verlag, New York, 1993.
- [7] Greg Friedman, *Intersection homology and Poincaré duality on homotopically stratified spaces*, submitted.
- [8] ———, *On the chain-level intersection pairing for PL pseudomanifolds*, submitted.
- [9] ———, *Stratified fibrations and the intersection homology of the regular neighborhoods of bottom strata*, *Topology Appl.* **134** (2003), 69–109.
- [10] ———, *Superperverse intersection cohomology: stratification (in)dependence*, *Math. Z.* **252** (2006), 49–70.
- [11] ———, *Singular chain intersection homology for traditional and super-perversities*, *Trans. Amer. Math. Soc.* **359** (2007), 1977–2019.
- [12] Mark Goresky, *Witt space cobordism theory*, *Intersection Cohomology* (et. al. A. Borel, ed.), *Progress in Mathematics*, vol. 50, Birkhauser, Boston, 1984, pp. 209–214.
- [13] Mark Goresky and Robert MacPherson, *Intersection homology theory*, *Topology* **19** (1980), 135–162.
- [14] ———, *Intersection homology II*, *Invent. Math.* **72** (1983), 77–129.
- [15] Mark Goresky and Paul Siegel, *Linking pairings on singular spaces*, *Comment. Math. Helvetici* **58** (1983), 96–110.
- [16] Daniel Huybrechts, *Complex geometry: An introduction*, Springer, New York, 2005.
- [17] Henry C. King, *Topological invariance of intersection homology without sheaves*, *Topology Appl.* **20** (1985), 149–160.
- [18] Frances Kirwan and Jonathan Woolf, *An introduction to intersection homology theory. second edition*, Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [19] T.Y. Lam, *The algebraic theory of quadratic forms*, W.A. Benjamin, Inc., Reading, Massachusetts, 1973.
- [20] Robert MacPherson and Kari Vilonen, *Elementary construction of perverse sheaves*, *Invent. Math.* **84** (1986), 403–435.
- [21] Clint McCrory, *Stratified general position*, *Algebraic and geometric topology* (Proc. Sympos., Univ. California, Santa Barbara, Calif. 1977) (Berlin), *Lecture Notes in Math.*, vol. 664, Springer, 1978, pp. 142–146.
- [22] J. Milnor and D. Husemoller, *Symmetric bilinear forms*, Springer Verlag, New York, 1973.

- [23] John W. Milnor and James D. Stasheff, *Characteristic classes*, Princeton University Press, Princeton, N. J., 1974, Annals of Mathematics Studies, No. 76. MR MR0440554 (55 #13428)
- [24] James R. Munkres, *Elements of algebraic topology*, Addison-Wesley, Reading, MA, 1984.
- [25] William L. Pardon, *Intersection homology Poincaré spaces and the characteristic variety theorem*, Comment. Math. Helv. **65** (1990), no. 2, 198–233.
- [26] Yuli B. Rudyak, *On Thom spectra, orientability, and cobordism*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998, With a foreword by Haynes Miller. MR MR1627486 (99f:55001)
- [27] P.H. Siegel, *Witt spaces: a geometric cycle theory for KO-homology at odd primes*, American J. Math. **110** (1934), 571–92.
- [28] Robert M. Switzer, *Algebraic topology - homology and homotopy*, Springer, Berlin, 2002.
- [29] Laurence Taylor and Bruce Williams, *Local surgery: foundations and applications*, Algebraic Topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978) (Berlin), Lecture Notes in Math., vol. 763, Springer, 1979, pp. 173–207.

Several diagrams in this paper were typeset using the \TeX commutative diagrams package by Paul Taylor.

Corrigendum to “Intersection homology with field coefficients: K -Witt spaces and K -Witt bordism”

Greg Friedman*

August 14, 2012

The author’s paper [2] concerns K -Witt spaces and, in particular, a computation of the bordism theory of such spaces. However, there is an error in the computation of the coefficient groups in dimensions $4k + 2$ when $\text{char}(K) = 2$. In this corrigendum, we state, as far as possible, the correct results. Details can be found in [1].

If we consider K -Witt spaces and K -Witt bordism using K -orientations, then for $\text{char}(K) = 2$, this is unoriented bordism, which we denote $\mathcal{N}_*^{K\text{-Witt}}$.

Theorem 1. *For a field K with $\text{char}(K) = 2$ and for¹ $i \geq 0$,*

$$\mathcal{N}_i^{K\text{-Witt}} \cong \begin{cases} \mathbb{Z}_2, & i \equiv 0 \pmod{2}, \\ 0, & i \equiv 1 \pmod{2}. \end{cases}$$

This result is also provided without detailed proof by Goresky in [3, page 498].

If we consider K -Witt spaces and K -Witt bordism using \mathbb{Z} -orientations, then we denote the bordism theory by $\Omega_*^{K\text{-Witt}}$. In this case, there remains one ambiguity in the computation, but we can show the following:

Theorem 2. *For a field K with $\text{char}(K) = 2$ and $k \geq 0$,*

1. $\Omega_0^{K\text{-Witt}} \cong \mathbb{Z}$,
2. $\Omega_{4k}^{K\text{-Witt}} \cong \mathbb{Z}_2$,
3. $\Omega_{4k+1}^{K\text{-Witt}} = \Omega_{4k+3}^{K\text{-Witt}} = 0$,
4. *Either*
 - (a) $\Omega_{4k+2}^{K\text{-Witt}} = 0$ for all k , or
 - (b) there exists some $N > 0$ such that $\Omega_{4k+2}^{K\text{-Witt}} = 0$ for all $k < N$ and $\Omega_{4k+2}^{K\text{-Witt}} \cong \mathbb{Z}_2$ for all $k \geq N$.

*This work was partially supported by a grant from the Simons Foundation (#209127 to Greg Friedman)

¹Since these are geometric bordism groups, they vanish in negative degree.

See [1] for a discussion of the difficulty in determining which case of item (4) of Theorem 2 holds.

Independent of the existence or value of N in condition (4) of the theorem, the computations from [2, Section 4.5] of $\Omega_*^{K\text{-Witt}}(\cdot)$ as a generalized homology theory on CW complexes continue to hold and to imply that for $\text{char}(K) = 2$,

$$\Omega_n^{K\text{-Witt}}(X) \cong \bigoplus_{r+s=n} H_r(X; \Omega_s^{K\text{-Witt}}).$$

Similarly,

$$\mathcal{N}_n^{K\text{-Witt}}(X) \cong \bigoplus_{r+s=n} H_r(X; \mathcal{N}_s^{K\text{-Witt}}).$$

Other minor errata. In [2] it should not be part of the definition of a K -Witt space that the space be irreducible as a pseudomanifold. However, as every K -Witt space of dimension > 0 is bordant to an irreducible K -Witt space [4, page 1099], this error does not affect the bordism group computations of [2]. Not every 0-dimensional K -Witt space is bordant to an irreducible one, but the computation of $\Omega_0^{K\text{-Witt}}$ reduces to the manifold theory and gives the result of [2] if one removes irreducibility from the definition.

The argument that $\Omega_{4k+2}^{K\text{-Witt}} = 0$ given in [2] does not hold when $k = 0$. However, in this dimension it is not difficult to prove the result directly; details are provided in [1].

References

- [1] Greg Friedman, *K-Witt bordism in characteristic 2*, preprint; see also <http://faculty.tcu.edu/gfriedman>.
- [2] ———, *Intersection homology with field coefficients: K-Witt spaces and K-Witt bordism*, *Comm. Pure Appl. Math.* **62** (2009), 1265–1292.
- [3] R. Mark Goresky, *Intersection homology operations*, *Comment. Math. Helv.* **59** (1984), no. 3, 485–505.
- [4] P.H. Siegel, *Witt spaces: a geometric cycle theory for KO-homology at odd primes*, *American J. Math.* **110** (1934), 571–92.

K-Witt bordism in characteristic 2

Greg Friedman*

August 13, 2012

Abstract

This note provides a computation of the bordism groups of K -Witt spaces for fields K with characteristic 2. We provide a complete computation for the unoriented bordism groups. For the oriented bordism groups, a nearly complete computation is provided as well a discussion of the difficulty of resolving a remaining ambiguity in dimensions equivalent to $2 \pmod{4}$. This corrects an error in the $\text{char}(K) = 2$ case of the author's prior computation of the bordism groups of K -Witt spaces for an arbitrary field K .

In [1], an n -dimensional K -Witt space, for a field K , is defined¹ to be an oriented compact n -dimensional PL stratified pseudomanifold X satisfying the K -Witt condition that the lower-middle perversity intersection homology group $I^{\bar{m}}H_k(L; K)$ is 0 for each link L^{2k} of each stratum of X of dimension $n - 2k - 1$, $k > 0$. Following the definition of stratified pseudomanifold in [2], X does not possess codimension one strata. Orientability is determined by the orientability of the top (regular) strata. This definition generalizes Siegel's definition in [11] of \mathbb{Q} -Witt spaces (called there simply "Witt spaces"). The motivation for this definition is that such spaces possess intersection homology Poincaré duality $I^{\bar{m}}H_i(X; K) \cong \text{Hom}(I^{\bar{m}}H_{n-i}(X; K), K)$.

The author's paper [1] concerns K -Witt spaces and, in particular, a computation of the bordism theory $\Omega_*^{K\text{-Witt}}$ of such spaces. However, there is an error in [1] in the computation of the coefficient groups $\Omega_{4k+2}^{K\text{-Witt}}$ when $\text{char}(K) = 2$.

It is claimed in [1] that $\Omega_{4k+2}^{K\text{-Witt}} = 0$. When $\text{char}(K) > 2$, the null-bordism of a $4k + 2$ dimensional K -Witt space X is established in [1] by following Siegel's computation [11] for \mathbb{Q} -Witt spaces by first performing a surgery to make the space irreducible and then performing

*This work was partially supported by a grant from the Simons Foundation (#209127 to Greg Friedman)

2000 Mathematics Subject Classification: 55N33, 57Q20, 57N80

Keywords: intersection homology, Witt bordism, Witt space

¹There is a minor error in [1] in that Witt spaces are stated to be irreducible, meaning that there is only a single top dimensional stratum. In general, this should not be part of the definition of a K -Witt space; cf. [11]. However, as every K -Witt space of dimension > 0 is bordant to an irreducible K -Witt space (see [11, page 1099]), this error does not affect the bordism group computations of [1]. It is not true that every 0-dimensional K -Witt space is bordant to an irreducible K -Witt space, but in this dimension the computations all reduce to the manifold theory and the computations given for this dimension in [1] are also correct if one removes irreducibility from the definition.

a sequence of singular surgeries to obtain a space X' such that $I^{\bar{m}}H_{2k+1}(X'; K) = 0$. The K -Witt null-bordism of X is the union of the trace of the surgeries from X to X' with the closed cone $\bar{c}X'$. One performs the singular surgeries on elements $[z] \in I^{\bar{m}}H_{2k+1}(X; K)$ such that $[z] \cdot [z] = 0$, where \cdot denotes the Goresky-MacPherson intersection product [2]. As the intersection product is skew symmetric on $I^{\bar{m}}H_{2k+1}(X; K)$, such a $[z]$ always exists. The error in [1] stems from overlooking that this last fact is not necessarily true in characteristic 2, where skew symmetric forms and symmetric forms are the same thing and so skew-symmetry does not imply $[z] \cdot [z] = 0$.

Corrected computations. To begin to remedy the error of [1], we first observe that it remains true in characteristic 2 that the map² $w : \Omega_{4k+2}^{\mathbb{Z}_2\text{-Witt}} \rightarrow W(\mathbb{Z}_2)$ is injective, where $W(\mathbb{Z}_2)$ is the Witt group of \mathbb{Z}_2 and w takes the bordism class $[X]$ to the class of the intersection form on $I^{\bar{m}}H_{2k+1}(X; \mathbb{Z}_2)$. For $k > 0$, this fact can be proven as it is proven for $w : \Omega_{4j}^{K\text{-Witt}} \rightarrow W(K)$, $j > 0$, in [1]: if one assumes that the intersection form on X represents 0 in $W(\mathbb{Z}_2)$ then the intersection form is split, in the language of [7]; see [7, Corollary III.1.6]. And so $I^{\bar{m}}H_{2k+1}(X; \mathbb{Z}_2)$ will possess an isotropic (self-annihilating) element by [7, Lemma I.6.3]. The surgery argument can then proceed³. As $W(\mathbb{Z}_2) \cong \mathbb{Z}_2$ (see [7, Lemma IV.1.5]), it follows that $\Omega_{4k+2}^{\mathbb{Z}_2\text{-Witt}}$ is either 0 or \mathbb{Z}_2 .

This argument does not hold for $4k+2 = 2$ as in this case the dimensions are not sufficient to guarantee that every middle-dimensional intersection homology class is representable by an irreducible element, which is necessary for the surgery argument; see [11, Lemma 2.2]. However, all 2-dimensional Witt spaces must have at worst isolated singularities, and so in particular such a space must have the form $X \cong (\amalg S_i) / \sim$, where the S_i are closed oriented surfaces and the relation \sim glues them together along various isolated points. But then X is bordant to $\amalg S_i$. This can be seen via a sequence of pinch bordisms as defined by Siegel [11, Section II] that pinch together the regular neighborhoods of sets of points of $\amalg S_i$. To see that the bordism is via a Witt space, it is only necessary to observe that the link of the interior cone point in each such pinch bordism will be a wedge of S^2 s, and it is easy to compute that $I^{\bar{m}}H_1(\vee_i S^2; K) = 0$ for any K . But now, since all closed oriented⁴ surfaces bound, $\Omega_2^{\mathbb{Z}_2\text{-Witt}} = 0$. This special case was also over-looked in [1], though this argument holds for any field K and is consistent with the claim of [1] that $\Omega_2^{K\text{-Witt}} = 0$ for all K .

Thus we have shown that $w : \Omega_{4k+2}^{\mathbb{Z}_2\text{-Witt}} \rightarrow W(\mathbb{Z}_2) \cong \mathbb{Z}_2$ is an injection for $k \geq 0$, trivially

²Recall from [1, Corollary 4.3] that the bordism groups depend only on the characteristic of the field, so for characteristic 2 it suffices to consider $K = \mathbb{Z}_2$.

³There is one other possible complication due to characteristic 2 that must be checked but that does not provide difficulty in the end: For characteristic not equal to 2, every split form is isomorphic to an orthogonal sum of hyperbolic planes [7, Lemma I.6.3], and this appears to be used in the proof of Theorem 4.4 of [11], which is heavily referenced in [1]. For characteristic 2, one can only conclude that a split form is isomorphic to one with matrix $\begin{pmatrix} 0 & I \\ I & A \end{pmatrix}$ for some matrix A . However, a detailed reading of the proof of [11, Theorem 4.4, particularly page 1097] reveals that it is sufficient to have a basis $\{\alpha, \beta, \gamma_1, \dots, \gamma_{2m}\}$ such that $\alpha \cdot \alpha = \alpha \cdot \gamma_i = 0$ for all i and $\alpha \cdot \beta = 1$, and this is certainly provided by a form with the given matrix.

⁴Recall that \mathbb{Z}_2 -Witt spaces are assumed to be \mathbb{Z} -oriented, though see below for more on orientation considerations

so for $k = 0$. Unfortunately, the question of surjectivity of w in dimensions $4k + 2$ is more complicated and not yet fully resolved. We can, however, make the following observation: if X is a \mathbb{Z}_2 -Witt space of dimension $4k - 2$, then⁵ $w([X \times \mathbb{C}P^2]) = w([X])$. So if there is a non-trivial element of $\Omega_{4k-2}^{\mathbb{Z}_2\text{-Witt}}$, then there is a non-trivial element of $\Omega_{4k+2}^{\mathbb{Z}_2\text{-Witt}}$.

Putting this together with the computations from [1] of $\Omega_*^{K\text{-Witt}}$ in dimension $\not\equiv 4k + 2 \pmod{4}$ (which remain correct), we have the following theorem:

Theorem 1. *For a field K with $\text{char}(K) = 2$, $\Omega_*^{K\text{-Witt}} = \Omega_*^{\mathbb{Z}_2\text{-Witt}}$, and for⁶ $k \geq 0$,*

1. $\Omega_0^{K\text{-Witt}} \cong \mathbb{Z}$,
2. for $k > 0$, $\Omega_{4k}^{K\text{-Witt}} \cong \mathbb{Z}_2$, generated by $[\mathbb{C}P^{2k}]$,
3. $\Omega_{4k+3}^{K\text{-Witt}} = \Omega_{4k+1}^{K\text{-Witt}} = 0$,
4. *Either*
 - (a) $\Omega_{4k+2}^{K\text{-Witt}} = 0$ for all k , or
 - (b) *there exists some $N > 0$ such that $\Omega_{4k+2}^{K\text{-Witt}} = 0$ for all $k < N$ and $\Omega_{4k+2}^{K\text{-Witt}} \cong \mathbb{Z}_2$ for all $k \geq N$.*

We will provide below some further discussion of the difficulties of deciding which case of (4) holds after discussing unoriented bordism.

Remark. Independent of the existence or value of N in condition (4) of the theorem, the computations from [1, Section 4.5] of $\Omega_*^{K\text{-Witt}}(\cdot)$ as a generalized homology theory on CW complexes continue to hold and to imply that for $\text{char}(K) = 2$,

$$\Omega_n^{K\text{-Witt}}(X) = \Omega_n^{\mathbb{Z}_2\text{-Witt}}(X) \cong \bigoplus_{r+s=n} H_r(X; \Omega_s^{\mathbb{Z}_2\text{-Witt}}).$$

Unoriented bordism. Given the motivation to recognize spaces that possess a form of Poincaré duality, it seems reasonable to consider K -Witt spaces that are K -oriented. This has no effect when $\text{char}(K) \neq 2$, in which case K -orientability is equivalent to \mathbb{Z} -orientability as considered in [1]. But when $\text{char}(K) = 2$, all pseudomanifolds are \mathbb{Z}_2 -orientable, which is equivalent to being K orientable, and the Poincaré duality isomorphism $I^{\bar{m}}H_k(X; K) \cong \text{Hom}(I^{\bar{m}}H_{n-k}(X; K), K)$ holds for all such compact pseudomanifolds satisfying the K -Witt condition.

If we allow K -Witt spaces and K -Witt bordism using K -orientations, then for $\text{char}(K) = 2$ we are essentially talking about unoriented bordism⁷, so to clarify the notation, let us denote the resulting bordism groups by $\mathcal{N}_*^{K\text{-Witt}}$. These groups can be computed as follows:

⁵Recall that the Künneth theorem holds within a single perversity when one term is a manifold, so we can compute the intersection forms of such product spaces in the usual way; see e.g. [6].

⁶Since these are geometric bordism groups, they vanish in negative degree.

⁷One could also define unoriented bordism groups of unoriented compact PL pseudomanifolds satisfying the K -Witt condition with $\text{char}(K) \neq 2$, but it is not clear how to study such groups by the present techniques, as there is no reason to expect that $I^{\bar{m}}H_*(X; K)$ would satisfy Poincaré duality for such a space X .

Theorem 2. For a field K with $\text{char}(K) = 2$ and for $i \geq 0$,

$$\mathcal{N}_i^{K\text{-Witt}} \cong \begin{cases} \mathbb{Z}_2, & i \equiv 0 \pmod{2}, \\ 0, & i \equiv 1 \pmod{2}. \end{cases}$$

Since writing [1], the author has discovered that this theorem is also provided without detailed proof by Goresky in [4, page 498]. We provide here the details:

Proof. It continues to hold that the local Witt condition depends only on the characteristic of K for the reasons provided in [1], so we may assume $K = \mathbb{Z}_2$. To see that $\mathcal{N}_n^{\mathbb{Z}_2\text{-Witt}} = 0$ for n odd, we simply note that X bounds the closed cone $\bar{c}X$, which is a \mathbb{Z}_2 -Witt space. The map $w : \mathcal{N}_{2k}^{\mathbb{Z}_2\text{-Witt}} \rightarrow W(\mathbb{Z}_2) \cong \mathbb{Z}_2$ is onto for each $k > 0$, as the intersection pairing on the \mathbb{Z}_2 -coefficient middle-dimensional homology of the real projective space $\mathbb{R}P^{2k}$ corresponds to the generator of $W(\mathbb{Z}_2)$ represented by the matrix $\langle 1 \rangle$. Furthermore, w is injective for $k > 1$ as in the preceding surgery argument, which does not rely on whether or not X is oriented, only on the existence of the intersection pairing over \mathbb{Z}_2 . In dimension 0, we have unoriented manifold bordism of points, so $\mathcal{N}_0^{\mathbb{Z}_2\text{-Witt}} \cong \mathbb{Z}_2$. Finally, as in the argument above for $\Omega_2^{\mathbb{Z}_2\text{-Witt}}$, the group $\mathcal{N}_2^{\mathbb{Z}_2\text{-Witt}}$ must be generated by closed surfaces (now not necessarily oriented), so $\mathcal{N}_2^{\mathbb{Z}_2\text{-Witt}}$ is a quotient of the unoriented manifold bordism group $\mathcal{N}_2 \cong \mathbb{Z}_2$; thus $\mathcal{N}_2^{\mathbb{Z}_2\text{-Witt}}$ must be isomorphic to \mathbb{Z}_2 as w maps $\mathbb{R}P^2$ onto the non-trivial element of $W(\mathbb{Z}_2) \cong \mathbb{Z}_2$. \square

Remark. An even simpler version of the argument of [1] implies that as a generalized homology theory

$$\mathcal{N}_n^{K\text{-Witt}}(X) \cong \bigoplus_{r+s=n} H_r(X; \mathcal{N}_s^{K\text{-Witt}})$$

for $\text{char}(K) = 2$, as in this case one no longer needs a separate argument to handle the odd torsion that can arise in $H_n(X; \Omega_0^{K\text{-Witt}})$ as a result of $\Omega_0^{K\text{-Witt}} \cong \mathbb{Z}$ not being 2-primary.

Further discussion of oriented bordism. We next provide some results that demonstrate the difficulty of determining which case of item (4) of Theorem 1 holds.

We will first see that $w([M]) = 0$ for any \mathbb{Z} -oriented manifold: Since dimension mod 2 is the only invariant⁸ of $W(\mathbb{Z}_2)$, this is a consequence of the following lemma, recalling that for a manifold, $I^{\bar{m}}H_*(M) = H_*(M)$.

Lemma. Let M be a closed connected \mathbb{Z} -oriented manifold of dimension $4k + 2$. Then $\dim(H_{2k+1}(M; \mathbb{Z}_2)) \equiv 0 \pmod{2}$.

Proof. By the universal coefficient theorem,

$$H_{2k+1}(M; \mathbb{Z}_2) \cong (H_{2k+1}(M) \otimes \mathbb{Z}_2) \oplus (H_{2k}(M) * \mathbb{Z}_2),$$

⁸As observed in the proof of [7, Lemma III.3.3], rank mod 2 yields a homomorphism $W(F) \rightarrow \mathbb{Z}_2$ for any field F . Since we know that $W(\mathbb{Z}_2) \cong \mathbb{Z}_2$ and that $\langle 1 \rangle$, which has rank 1, is a generator of $W(F)$ (it is certainly non-zero, using [7, Lemma I.6.3 and Lemma III.1.6]), it follows that rank mod 2 determines the isomorphism.

where the asterisk denotes the torsion product. Let $T_*(M)$ denote the torsion subgroup of $H_*(M)$, and let $T_*^2(M)$ denote $T_*(M) \otimes \mathbb{Z}_2 \cong T_*(M) * \mathbb{Z}_2$; the isomorphism follows from basic homological algebra because $T_*(M)$ is a finite abelian group. $T_*^2(M)$ is a direct sum of \mathbb{Z}_2 terms. Then $H_{2k+1}(M) \otimes \mathbb{Z}_2 \cong \mathbb{Z}_2^B \oplus T_{2k+1}^2(M)$, where B is the $2k + 1$ Betti number of M , and $H_{2k}(M) * \mathbb{Z}_2 \cong T_{2k}^2(M)$. Thus $H_{2k+1}(M; \mathbb{Z}_2) \cong \mathbb{Z}_2^B \oplus T_{2k+1}^2(M) \oplus T_{2k}^2(M)$. Since M is a closed \mathbb{Z} -oriented manifold, there is a nondegenerate skew-symmetric intersection form on $H_{2k+1}(M; \mathbb{Q})$, and so B is even. Since M is a closed \mathbb{Z} -oriented manifold, the nonsingular linking pairing $T_{2k+1}(M) \otimes T_{2k}(M) \rightarrow \mathbb{Q}/\mathbb{Z}$ gives rise to an isomorphism $T_{2k+1}(M) \cong \text{Hom}(T_{2k}(M), \mathbb{Q}/\mathbb{Z})$, and since $\text{Hom}(\mathbb{Z}_n, \mathbb{Q}/\mathbb{Z}) \cong \mathbb{Z}_n$, it follows that $T_{2k+1}(M) \cong T_{2k}(M)$. Therefore $T_{2k+1}^2(M) \cong T_{2k}^2(M)$. Thus $H_{2k+1}(M; \mathbb{Z}/2)$ consists of an even number of \mathbb{Z}_2 terms. \square

Remark. Since the lemma utilizes only integral Poincaré duality and the universal coefficient theorem, it follows that, in fact, $w([X]) = 0$ for any IP space⁹; these are spaces that satisfy local conditions guaranteeing that intersection homology Poincaré duality holds over the integers and that a universal coefficient theorem holds (see [3, 10]).

A slightly more elaborate argument demonstrates that it is also not possible to have $w([X]) \neq 0$ if X is a \mathbb{Z} -oriented \mathbb{Z}_2 -Witt space with at worst isolated singularities:

Proposition. *Let X be a closed \mathbb{Z} -oriented $4k + 2$ -dimensional \mathbb{Z}_2 -Witt space with at worst isolated singularities. Then $w([X]) = 0$.*

Proof. Since X has at worst point singularities, it follows from basic intersection homology calculations (see [2, Section 6.1]) that $I^{\bar{m}}H_{2k+1}(X; \mathbb{Z}_2) \cong \text{im}(H_{2k+1}(M; \mathbb{Z}_2) \rightarrow H_{2k+1}(M, \partial M; \mathbb{Z}_2))$, where M is the compact \mathbb{Z} -oriented PL ∂ -manifold obtained by removing an open regular neighborhood of the singular set of X . We will show that if $[z] \in \text{im}(H_{2k+1}(M; \mathbb{Z}_2) \rightarrow H_{2k+1}(M, \partial M; \mathbb{Z}_2))$, then the intersection product $[z] \cdot [z] = 0$. It follows that the intersection pairing on $I^{\bar{m}}H_{2k+1}(X; \mathbb{Z}_2)$ is split by [7, Lemma III.1.1], since then there can be no non-trivial anisotropic subspace. This implies that $w([X]) = 0$ by the definition of the Witt group.

The following argument that $[z] \cdot [z] = 0$ was suggested by “Martin O” on the web site MathOverflow [9]. By Poincaré duality, it suffices to show that $\alpha \cup \alpha = 0$, where α is the Poincaré dual of $[z]$ in $H^{2k+1}(M, \partial M; \mathbb{Z}_2)$. But now $\alpha \cup \alpha = Sq^{2k+1}\alpha = Sq^1Sq^{2k}\alpha = \beta^*Sq^{2k}\alpha$, where β^* is the Bockstein associated with the sequence $0 \rightarrow \mathbb{Z}_2 \rightarrow \mathbb{Z}_4 \rightarrow \mathbb{Z}_2 \rightarrow 0$ (see [5, Section 4.L]). In the case at hand, this is the Bockstein $\beta^* : H^{4k+1}(M, \partial M; \mathbb{Z}_2) \rightarrow H^{4k+2}(M, \partial M; \mathbb{Z}_2)$. But this map is trivial. To see this, observe that there is a commutative

⁹Also called “intersection homology Poincaré spaces,” though this is perhaps a misnomer as “Poincaré spaces” are generally not required to be manifolds while IP spaces are still expected to be pseudomanifolds.

diagram

$$\begin{array}{ccc}
 H^{4k+1}(M, \partial M; \mathbb{Z}_2) & \xrightarrow{\beta^*} & H^{4k+2}(M, \partial M; \mathbb{Z}_2) \\
 \downarrow \cong & & \downarrow \cong \\
 H_1(M; \mathbb{Z}_2) & \xrightarrow{\beta_*} & H_0(M; \mathbb{Z}_2),
 \end{array}$$

where β_* is the homology Bockstein and the vertical maps are Poincaré duality. The existence of this diagram follows as in [8, Lemma 69.2]. But now $\beta_* : H_1(M; \mathbb{Z}_2) \rightarrow H_0(M; \mathbb{Z}_2)$ is trivial, as the standard map $\times 2 : H_0(M; \mathbb{Z}_2) \rightarrow H_0(M; \mathbb{Z}_4)$ is injective. \square

Hence any candidate to have $w([X]) = 1$ must have singular set of dimension > 0 and must not be an IP space. Given that all K -Witt spaces for $\text{char}(K) \neq 2$ are K -Witt bordant to spaces with at worst isolated singularities [11, 1], it is unclear how to proceed to determine whether \mathbb{Z}_2 -Witt spaces with $w([X]) = 1$ exist. One method to prove that they do not would be to try to show “by hand” that every \mathbb{Z}_2 -Witt space is \mathbb{Z}_2 -Witt bordant to a space with at most isolated singularities, but the only proof currently known to the author of this fact for fields of other characteristics utilizes the bordism computations of [11, 1].

References

- [1] Greg Friedman, *Intersection homology with field coefficients: K -Witt spaces and K -Witt bordism*, Comm. Pure Appl. Math. **62** (2009), 1265–1292.
- [2] Mark Goresky and Robert MacPherson, *Intersection homology theory*, Topology **19** (1980), 135–162.
- [3] Mark Goresky and Paul Siegel, *Linking pairings on singular spaces*, Comment. Math. Helvetici **58** (1983), 96–110.
- [4] R. Mark Goresky, *Intersection homology operations*, Comment. Math. Helv. **59** (1984), no. 3, 485–505.
- [5] Allen Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.
- [6] Henry C. King, *Topological invariance of intersection homology without sheaves*, Topology Appl. **20** (1985), 149–160.
- [7] J. Milnor and D. Husemoller, *Symmetric bilinear forms*, Springer Verlag, New York, 1973.
- [8] James R. Munkres, *Elements of algebraic topology*, Addison-Wesley, Reading, MA, 1984.
- [9] Martin O, see <http://mathoverflow.net/questions/53419/>.

- [10] William L. Pardon, *Intersection homology Poincaré spaces and the characteristic variety theorem*, Comment. Math. Helvetici **65** (1990), 198–233.
- [11] P.H. Siegel, *Witt spaces: a geometric cycle theory for KO-homology at odd primes*, American J. Math. **110** (1934), 571–92.