

Quasi-representations of Surface Groups

José R. Carrión

Purdue University

joint work with Marius Dadarlat

East Coast Operator Algebras Symposium

University of Tennessee, Knoxville

October 6, 2012

- Idea: obtain numerical invariants by pushing-forward K -theory elements via *approximately* multiplicative maps.

$$\pi: B \rightarrow M_n(\mathbb{C}) \rightsquigarrow \pi_{\#}: K_0(B) \rightarrow \mathbb{Z}$$

Why *approximately* multiplicative?

Proposition (Dadarlat)

Suppose Γ satisfies Baum-Connes and is torsion-free. If

$$\pi: C^*(\Gamma) \rightarrow M_n(\mathbb{C})$$

is a (unital) representation, then the induced map π_* on K_0 equals $n \cdot \iota_*$ where $\iota =$ trivial rep. of Γ .

Definition of a quasi-representation

$A =$ unital C^* -algebra (with a tracial state τ)

Definition

Let $\mathcal{F} \subset \Gamma$ be finite, $\varepsilon > 0$.

$\pi: \Gamma \rightarrow U(A)$ is an $(\mathcal{F}, \varepsilon)$ -representation if $\forall s, t \in \mathcal{F}$:

- $\pi(1) = 1_A$
- $\|\pi(s^{-1}) - \pi(s)^*\| < \varepsilon$
- $\|\pi(st) - \pi(s)\pi(t)\| < \varepsilon$

- *Quasi-representation*: \mathcal{F} , ε not necessarily specified.
- May extend a quasi-rep π of Γ to a unital, linear, *approximately multiplicative* contraction on $\ell^1(\Gamma)$ (in the obvious way).

Pushing-forward via quasi-representations

Let $\pi: \Gamma \rightarrow U(A)$ be a quasi-representation. (Extend to $\ell^1(\Gamma)$.)

How to push-forward $x \in K_0(\ell^1(\Gamma))$

- choose idempotents e_0, e_1 in matrices over $\ell^1(\Gamma)$ s.t. $x = [e_0] - [e_1]$.
- π multiplicative enough $\Rightarrow \pi(e_i) \approx$ idempotent in matrices over A
 $\Rightarrow \pi(e_i)$ may be perturbed to idempotent f_i over A
- define $\pi_{\#}(x) = [f_0] - [f_1] \in K_0(A)$.

To obtain numerical invariants, apply trace $\tau \in T(A)$.

Example: Almost commuting unitaries

Suppose $u, v \in U(n)$ and $\|uvu^{-1}v^{-1} - 1_n\|$ is small.

- Associate a quasi-representation $\pi: \mathbb{Z}^2 \rightarrow U(n)$ s.t.

$$s \mapsto u, \quad t \mapsto v, \quad st \mapsto uv$$

- $K_0(\ell^1(\mathbb{Z}^2)) \cong K_0(C(\mathbb{T}^2)) = \mathbb{Z}[1] \oplus \mathbb{Z}\beta$. ($\beta =$ Bott element)
- $\|[u, v] - 1_n\|$ small enough $\Rightarrow \pi$ multiplicative enough so that

$$\kappa(u, v) := \text{Tr } \pi_{\#}(\beta) \in \mathbb{Z}$$

is defined.

Theorem (Exel-Loring '91)

If $\|[u, v] - 1_n\|$ is small enough, then

$$\kappa(u, v) = \frac{1}{2\pi i} \text{Tr } \log([u, v]).$$

Setup for main result

Can we generalize Exel-Loring? In what context?

M = g -holed torus (for some $g \in \mathbb{N}$)

$\Gamma := \pi_1(M)$. Known that:

$$\Gamma = \left\langle s_1, t_1, \dots, s_g, t_g : \prod_{i=1}^g [s_i, t_i] \right\rangle$$

$$([s_i, t_i] := s_i t_i s_i^{-1} t_i^{-1})$$

A = unital C^* -algebra with a tracial state τ

Fix a quasi-representation $\pi: \Gamma \rightarrow U(A)$.

The analogue of the invariant $\kappa(u, v)$

$$\begin{array}{ccc}
 [M] & & \mathbb{Z} \oplus \mathbb{Z}\mu[M] \\
 \cap & & \parallel \\
 K_0(M) & \xrightarrow{\mu} & K_0(\ell^1(\Gamma)) \ni \mu[M] \\
 & & \downarrow \pi_{\#} \\
 & & K_0(A) \\
 & & \downarrow \tau \\
 & & \mathbb{R} \ni \tau(\pi_{\#}(\mu[M]))
 \end{array}$$

(Lafforgue's ℓ^1 -version of) the assembly map μ is an isomorphism in this case.

Theorem (C-Dadarlat)

Fix a genus $g \in \mathbb{N}$. There exist $\varepsilon > 0$ and a finite subset $\mathcal{F} \subset \Gamma$ s.t. the following holds:

If A is a unital C^* -algebra, $\tau \in T(A)$ and $\pi: \Gamma \rightarrow U(A)$ is any $(\mathcal{F}, \varepsilon)$ -representation, then

$$\tau(\pi_{\#}(\mu[M])) = \frac{1}{2\pi i} \tau \left(\log \left(\prod_{i=1}^g [\pi(s_i), \pi(t_i)] \right) \right).$$

Remark

$\forall \varepsilon > 0$, finite $\mathcal{F} \subset \Gamma \exists \delta > 0$ s.t.
given A and $u_1, v_1, \dots, u_g, v_g \in U(A)$ satisfying

$$\left\| \prod_{i=1}^g [u_i, v_i] - 1 \right\| < \delta,$$

then there is an $(\mathcal{F}, \varepsilon)$ -representation $\pi: \Gamma \rightarrow U(A)$ with $\pi(s_i) = u_i$ and $\pi(t_i) = v_i$.

Example: NC tori

$A_\theta := C^*(u, v \mid u, v \text{ are unitaries s.t. } vu = e^{2\pi i\theta} uv)$.

θ is small enough $\Rightarrow \exists$ quasi-representation $\pi: \mathbb{Z}^2 \rightarrow U(A_\theta)$ s.t.
 $\pi(s) = u, \pi(t) = v$ and

$$\tau(\pi_{\#}(\beta)) = \frac{1}{2\pi i} \tau(\log [u, v]) = -\theta.$$

The context for the formula

Recall the formula:

$$\tau(\pi_{\sharp}(\mu[M])) = \frac{1}{2\pi i} \tau \left(\log \left(\prod_{i=1}^g [\pi(s_i), \pi(t_i)] \right) \right)$$

The assembly map $\mu: K_0(M) \rightarrow K_0(\ell^1(\Gamma))$ is implemented by the *Mishchenko line bundle* ℓ :

$$\begin{array}{ccccc} K_0(C(M) \otimes \ell^1(\Gamma)) & & KK(C(M), \mathbb{C}) & & KK(\mathbb{C}, \ell^1(\Gamma)) \\ \psi & & \parallel & & \parallel \\ \ell & \times & K_0(M) & \xrightarrow{\mu} & K_0(\ell^1(\Gamma)) \end{array}$$

The context for the formula (cont.)

$$\begin{array}{ccccc}
 & & [M] \dashv \longrightarrow & & \mu[M] \\
 K_0(C(M) \otimes \ell^1(\Gamma)) \ni \ell & \times & K_0(M) \xrightarrow{\mu} & K_0(\ell^1(\Gamma)) & \\
 \downarrow (\text{id}_{C(M)} \otimes \pi)_\# & & \downarrow \text{ch} & \downarrow \pi_\# & \downarrow \\
 K_0(C(M) \otimes A) \ni \ell_\pi & & & K_0(A) & \\
 \downarrow \text{ch}_\tau & & & \downarrow \tau & \\
 H^{\text{even}}(M) \ni \text{ch}_\tau(\ell_\pi) \times H^{\text{even}}(M) & \longrightarrow & \mathbb{R} & & \tau(\pi_\#(\mu[M])) \\
 & & \langle \text{ch}_\tau(\ell_\pi), [M] \rangle & &
 \end{array}$$

Special case of a Theorem of Dadarlat:

$$\tau(\pi_\#(\mu[M])) = \langle \text{ch}_\tau(\ell_\pi), [M] \rangle$$

Hilbert A -module bundles

- $l \in K_0(C(M) \otimes \ell^1(\Gamma)) \Rightarrow$ push-forward $l_\pi \in K_0(C(M) \otimes A)$
- $K_0(C(M) \otimes A) =$ Grothendieck group of isomorphisms classes of f.g.p. *Hilbert A -module bundles*

Hilbert A -module bundle $E \rightarrow M$


- $E \rightarrow M$: fibers \cong Hilbert A -modules
- $E =$ f.g.p. $\Rightarrow E$ has (unique) *smooth structure*; curvature Ω
- For the proof of the theorem, we construct bundle $E_\pi \rightarrow M$ s.t. $[E_\pi] = l_\pi$.
- Construction is explicit enough that we may apply Chern-Weil theory, using $\text{ch}_\tau(l_\pi) = \tau(i\Omega/2\pi) \in \Omega^2(M, \mathbb{C})$, to get

$$\langle \text{ch}_\tau(l_\pi), [M] \rangle = \int_M \tau \left(\frac{i\Omega}{2\pi} \right).$$

Simplicial complexes and evaluation of the integral


- To construct E_π and deal with $\int_M \tau(i\Omega/2\pi)$ we work with a triangulation of M .
- Edges in simplicial complex \rightsquigarrow elements of $\Gamma = \pi_1(M)$

$g = 1$ case: \mathbb{T}^2



Evaluation of the integral (cont.)

On a 2-simplex $\sigma = \langle i, j, k \rangle \dots$



- $s_{ij} :=$ element of Γ corresp. to edge ij
- have “cocycle condition”: $s_{ij}s_{jk} = s_{ik}$
- π quasi-representation
 $\Rightarrow \pi(s_{ij})\pi(s_{jk}) \approx \pi(s_{ik})$
- in the \mathbb{T}^2 case:
 $\pi(t)\pi(s) = vu \approx uv = \pi(ts)$

Let $\xi_\sigma = \text{segment } \pi(s_{ij})\pi(s_{jk}) \rightsquigarrow \pi(s_{ik})$ in $GL_\infty(A)$. Then

$$\int_M \tau(\Omega) = \sum_\sigma (-1)^{\text{ord}(\sigma)} \tilde{\Delta}_\tau(\xi_\sigma).$$

where $\tilde{\Delta}_\tau =$ de la Harpe-Skandalis determinant.

To summarize

- Interested in how a quasi-rep π acts on $K_0(\ell^1(\Gamma))$, where $\Gamma =$ surface group
- $K_0(\ell^1(\Gamma)) \cong \mathbb{Z} \oplus \mathbb{Z}\mu[M]$

- Push-forward $\mu[M]$ via π and apply trace $\tau : \tau(\pi_{\#}(\mu[M]))$
- Push-forward ℓ , apply ch_{τ} etc. : $\langle \text{ch}_{\tau}(\ell_{\pi}), [M] \rangle$.

Dadarlat: red invariant = blue invariant.

- Use Chern-Weil theory for Hilbert A -module bundles to deal with $\text{ch}_{\tau}(\ell_{\pi})$.