



The spectrum of basic Dirac operators

기본 Dirac 사업자의 스펙트럼

Ken Richardson

켄 리차드슨

Jeju National University, Korea

제주대학교, 대한민국

August 12, 2009

This talk includes joint work with
Georges Habib, *A brief note on the
spectrum of the basic Dirac operator*, Bull.
London Math. Soc. **41** (2009), 683-690.

This talk also mentions joint work with

- Jochen Brüning
- Seoung Dal Jung
- Franz Kamber
- Jeff Lee
- Efton Park (*not Korean !!!*)



Plan of the talk

- Gomapsumida to the actual organizers
- Smooth foliations and basic cohomology
- Riemannian foliations
- Additional structure: bundle-like metrics
- The standard basic Laplacian
- Twisted duality for basic cohomology
- Definition of ordinary Dirac operator
- Definition of basic Dirac operators
- Invariance of the spectrum of basic Dirac operators
- A modified differential
- Poincare duality for modified basic cohomology

Smooth foliations

- Let (M, F) be a smooth manifold along with a smooth foliation F . This means that F is a partition of M into immersed submanifolds (*leaves*) such that the transition functions for the local product neighborhoods are smooth.
- Let TF denote the tangent bundle to the foliation. At each point p of M , $T_p F$ is the tangent space to the leaf through p .

Basic forms

- Basic forms of (M, F) are forms that are smooth on M but that locally are forms which depend only on the variables of the leaf spaces, the quotient of the local product by the leaves. Let $\Omega(M, F) \subset \Omega(M)$ denote the space of basic forms.

- Invariant definition:

$$\Omega(M, F) = \{ \beta \in \Omega(M) : i(X)\beta = 0,$$

$$i(X) d\beta = 0, \forall X \in TF \}$$

Basic cohomology

- Note that the exterior derivative ***d*** maps basic forms to basic forms, and thus we may define the basic cohomology as

with
$$H^k(M, F) = \frac{\ker d_k}{\operatorname{Im} d_{k-1}}$$

$$d_k = d : \Omega^k(M, F) \rightarrow \Omega^{k+1}(M, F)$$

- Physics-y interpretation: we are doing calculus with the set of leaves (strings?). One can think of basic forms as forms on the space of leaves (a singular, possibly non-Hausdorff space) or of leaf closures, and basic cohomology is a smooth version of the cohomology of that space. But not really.

Basic cohomology, continued

- Basic cohomology can be infinite dimensional. It can be really trivial.
- We can also define basic cohomology of forms twisted by a vector bundle. If one does this, one can learn a lot about the topology of the leaf space of the foliation.
- Basic Cohomology does not satisfy Poincaré duality. If there are additional restrictions, it may satisfy duality --- for example, if the manifold has a metric for which the leaves are equidistant and are minimal submanifolds.

Riemannian foliations

- Riemannian foliations are smooth foliations endowed with a holonomy-invariant metric on the quotient bundle $Q = TM/TF$.
- This means that Q has a metric g_Q such that
$$\theta(X)g_Q = 0, \quad \forall X \in TF$$

($\theta(X)$ means Lie derivative with respect to X .)
- One thinks of this as a metric on the (singular) space of leaves. But not really.

Additional structure: bundle-like metrics

- A *bundle-like metric* on a Riemannian foliation (M, F, g_Q) is a metric on M such that its restriction to $NF = (TF)^\perp$ agrees with g_Q through the natural isomorphism $NF \cong Q$.
- Every Riemannian foliation admits bundle-like metrics, and there are many choices of bundle-like metrics that are compatible with a given (M, F, g_Q) structure. One may freely choose the metric on the leaves and also the transverse subbundle NF .

Bundle-like metrics, continued

- A bundle-like metric on a smooth foliation is exactly a metric on the manifold such that the leaves of the foliation are locally equidistant.
- There are topological restrictions to the existence of bundle-like metrics (and thus Riemannian foliations). Important examples include:
 - The leaf closures partition the manifold (are disjoint).
 - The basic cohomology is finite-dimensional.
 - The orthogonal projection
$$P : L^2(\Omega(M)) \rightarrow L^2(\Omega(M, F))$$
maps the subspace of smooth forms onto the subspace of smooth basic forms.

Riemannian foliations, continued

Historical note:

Riemannian foliations were introduced by B. Reinhart in 1959. Good references for Riemannian foliations and bundle-like metrics can be found in the books and papers by B. Reinhart, F. Kamber, Ph. Tondeur, P. Molino.



The basic Laplacian

- On a Riemannian foliation with bundle-like metric, one can define a basic Laplacian that maps basic forms to basic forms:

$$\Delta_b = d\delta_b + \delta_b d : \Omega(M, F) \rightarrow \Omega(M, F)$$

where δ_b is the L^2 -adjoint of the restriction of d to basic forms: $\delta_b = P\delta$ is the ordinary adjoint of d followed by the orthogonal projection to the space of basic forms.

- This operator and its spectrum depend on the choice of bundle-like metric.

Basic Laplacian, continued

- The basic Laplacian can be used to define transversal heat flow on Riemannian foliations with bundle-like metrics. Such a flow corresponds to assuming that the leaves of the foliation are perfect conductors of heat.
- It turns out that the basic Laplacian is the restriction of a non-symmetric second order elliptic operator on all forms. Only in special cases is it the same as the ordinary Laplacian.
- It is also not the same as the formal Laplacian defined on the local quotient (or transversal). This transversal Laplacian is in general not symmetric on the space of basic forms, but it does preserve the basic forms.
- The basic heat flow asymptotics are more complicated than that of the standard heat kernel, but there is a fair amount known (Transactions AMS paper to appear).


The basic adjoint δ_b

Properties of the adjoint δ_b of the restriction of d to basic forms (see Park-R paper in Amer J Math):

$$\begin{aligned}\delta_b &= P\delta \\ &= \pm \overline{*} d \overline{*} + \kappa_b \lrcorner \\ &= \delta_T + \kappa_b \lrcorner,\end{aligned}$$

Here, $\overline{*}$ is the pointwise transversal Hodge star operator, and $\kappa_b \lrcorner$ denotes interior product with the projection of the mean curvature one-form onto basic forms, and δ_T is the formal adjoint of d on the local transversals.

The mean curvature form

- Let $H = \sum_{i=1}^p \pi(\nabla_{f_i}^M f_i)$ where $\pi : TM \rightarrow NF$ is the bundle projection, and $\{f_i\}_{i=1, \dots, p}$ is a local orthonormal frame of TF . This is the mean curvature vector field, and its dual one-form is $\kappa = H^\flat$. The projection $\kappa_b = P\kappa$ is the basic one-form on the previous slide.
- The operator $\kappa_b \lrcorner$ is the (pointwise) adjoint of the operator $\kappa_b \wedge$. Clearly, it depends on the choice of bundle-like metric.
- $\kappa_b = P\kappa$ is a closed form whose cohomology class in $H^1(M, F)$ is independent of the choice of bundle-like metric (see Álvarez-López )

Twisted duality for basic cohomology

- Even for Riemannian foliations, Poincaré duality does not hold for basic cohomology.
- However, note that $d - \kappa_b \wedge$ is also a differential which defines a cohomology of basic forms, and $\overline{*}\delta_b = \pm(d - \kappa_b \wedge)\overline{*}$, etc.
- Conclusion (Kamber/Tondeur/Park/R): if q is the codimension of the Riemannian foliation, then
$$H_d^*(M, F) \cong H_{d-\kappa_b \wedge}^{q-*}(M, F)$$

The ordinary Dirac-type operator


- Euclidean space:

$$D = \sum e_i \cdot \frac{\partial}{\partial x_i}, \quad e_i \cdot e_j + e_j \cdot e_i = -2\delta_{ij} \mathbf{1}$$

- 3 dimensions:

$$e_1 \cdot = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad e_2 \cdot = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad e_3 \cdot = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

- In general: $D = \sum e_i \cdot \nabla_i$
- Physics: eigensections correspond to particles with spin at different energy levels.



Examples of Dirac operators



- “The” spin-c Dirac operator
- The de Rham operator

$$d + \delta : \Omega^{\text{even}}(M) \rightarrow \Omega^{\text{odd}}(M)$$

- The signature operator

$$d + \delta : \Omega^+(M) \rightarrow \Omega^-(M)$$

Here, $\star = \pm *$, $\star^2 = 1$

$$\Omega^\pm(M) = \{\alpha \in \Omega^*(M) : \star \alpha = \pm \alpha\}$$

- The Dolbeault operator

$$\partial + \bar{\partial} : \Omega^{0,\text{even}}(M) \rightarrow \Omega^{0,\text{odd}}(M)$$

More on the de Rham operator

- The Laplacian is $(d + \delta)^2 = \Delta$

- We have

$$\ker(d + \delta) = \{ \text{harmonic forms} \}$$

- And

$$\begin{aligned} & \text{index}(d + \delta)|_{\Omega^{\text{even}}} \\ &= \dim \ker(d + \delta)|_{\Omega^{\text{even}}} - \dim \ker((d + \delta)|_{\Omega^{\text{even}}})^* \\ &= \chi(M) \end{aligned}$$

Basic Dirac operators

- Given a Riemannian foliation (M, F) of codim q with compatible bundle-like metric, E a foliated vector bundle,

$$D_{\text{tr}} s = \sum_{i=1}^q e_i \cdot \nabla_{e_i}^E s ,$$

$$D_b s = \frac{1}{2}(D_{\text{tr}} + D_{\text{tr}}^*)s = \sum_{i=1}^q e_i \cdot \nabla_{e_i}^E s - \frac{1}{2}\kappa_b^\# \cdot s$$

- References: Glazebrook-Kamber, S.D.Jung



Example: basic de Rham operator

- We have

$$D_{\text{tr}} = d + \delta_T = d + \delta_b - \kappa_b \lrcorner : \Omega^{\text{even}}(M, F) \rightarrow \Omega^{\text{odd}}(M, F)$$

$$D_b = \frac{1}{2}(D_{\text{tr}} + D_{\text{tr}}^*)s = d + \delta_b - \kappa_b \lrcorner - \frac{1}{2}\kappa_b^\# \cdot$$

$$= d + \delta_b - \kappa_b \lrcorner - \frac{1}{2}(\kappa_b \wedge - \kappa_b \lrcorner)$$

$$= d + \delta_b - \frac{1}{2}\kappa_b \lrcorner - \frac{1}{2}\kappa_b \wedge.$$

The spectrum of a basic Dirac operator

Theorem (Habib-R) Fix a Riemannian foliation. For any bundle-like metric and basic Dirac operator compatible with the Riemannian structure, the spectrum of this basic Dirac operator is independent of the choice of bundle-like metric.

a Riemannian foliation 수정합니다. 어떤 번들로 들어 - 미터법과 기본처럼 Dirac 연산자는 Riemannian 구조,이 기본 Dirac 연산자의 스펙트럼과 호환 번들의 선택과 독립적입니다 - 미터법처럼.

Idea of proof

- One can show that every different choice of bundle-like metric changes the L^2 -inner product by multiplication by a specific smooth, positive basic function.
- This changes the basic Dirac operator by a zeroth order operator that is Clifford multiplication by an exact basic one-form.
- This new operator is conjugate to the original one.

Consequences

- D. Dominguez showed that every Riemannian foliation admits a bundle-like metric for which the mean curvature form is basic.
- Further, March/MinOo/Ruh and A. Mason showed that the bundle-like metric may be chosen so that the mean curvature is basic-harmonic for that metric.
- Therefore, in calculating or estimating the eigenvalues of the basic Dirac operator, one may choose the bundle-like metric so that the mean curvature is basic-harmonic.
- Immediately we may obtain stronger inequalities for eigenvalue estimates.

A modified differential

- Recall the basic de Rham operator is

$$\begin{aligned} D_b &= d + \delta_b - \frac{1}{2}\kappa_b \lrcorner - \frac{1}{2}\kappa_b \wedge \\ &= \tilde{d} + \tilde{\delta} \end{aligned}$$

where $\tilde{d} = d - \frac{1}{2}\kappa_b \wedge$, $\tilde{\delta} = \delta_b - \frac{1}{2}\kappa_b \lrcorner$

- These are differentials and codifferentials on basic forms.
- Bonus: $\tilde{\delta} \overline{*} = \pm \overline{*} \tilde{d} \quad \dots \text{etc.}$
- Thus the basic signature operator can be defined!

New basic Laplacian and basic cohomology

- Let $\tilde{\Delta} = (\tilde{d} + \tilde{\delta})^2$. Then the eigenvalues of this operator depend only on the Riemannian foliation structure.
- The new basic cohomology satisfies Poincaré duality and basic Hodge theory, and the isomorphism is:
$$\overline{*} : H_{\tilde{d}}^*(M, F) \xrightarrow{\cong} H_{\tilde{d}}^{q-*}(M, F)$$
- Even though the differential depends on the choice of bundle-like metric, the cohomology groups are independent of that choice.

2009 NIMS International Workshop on

Differential Geometry and Related Topics

2009.8.10 ~ 2009.8.14 (International Center)

National Institute for Mathematical Sciences Sponsors : Jenu National University, KOSEF, KRF

Thank you for your
attention....